

CRD 92 Powder for Oral Solution

**Composition:** Each 100 gm of CRD 92 contains spiramycin (as adipate salt) 50 MIU, trimethoprim 5.0g.

**Product Description:** White powder with sugar taste.

**Pharmacodynamics:** The spiramycin is a bacteriostatic antibiotic belonging to the macrolides family. Spiramycin acts on bacterial protein synthesis by binding reversibly to bacterial 50 S ribosomal sub-unit. It binds to the donor site and inhibits translocation needed for peptide chain growth. This mechanism leads to an inhibition of the bacterial protein synthesis and consequently the growth of bacterial culture is stopped. Spiramycin acts essentially on organisms with rapid division. The activity of spiramycin is mainly bacteriostatic.

Cross-resistance within the macrolide group of antibiotics cannot be excluded. Such resistance is generally from plasmid origin in streptococcus and staphylococcus.

Trimethoprim is an anti-infective agent belonging to diaminopyrimidine group. It exerts its bactericidal activity by inhibiting the folic acid biosynthesis. It inhibits bacterial dihydrofolate reductase.

The activity spectrum of this combination includes Gram-positive organisms (*Staphylococcus*, ...) and Gram-negative organisms (*Echerichia coli*, *Salmonella*, *Proteus*, *Enterobacter*, *Bordetella*...) and mycoplasma (*Mycoplasma spp*).

**Pharmacokinetics:** Following oral administration, spiramycin is rapidly absorbed and diffuses in all tissues and serous except meninge. The plasma protein binding is very weak and the tissues exhibiting the greatest concentration are the lungs, spleen, kidneys, liver and bones. Spiramycin passes through the placental barrier and is found in maternal milk.

Spiramycin is mainly excreted by biliary route but an important proportion is reabsorbed by small intestine (enterohepatic cycle). Spiramycin is excreted by faeces essentially as active neospiramycin. Trimethoprim is rapidly absorbed after oral administration. Diffusion occurs into the whole organism. It is partly metabolised into liver. Its elimination is mainly renal.

**Indication:** Calves, pigs and poultry: treatment of generalised, respiratory and digestive infections due to bacteria susceptible to spiramycin and trimethoprim.

**Routes of administration:** Oral route in milk, lactating feed, liquid feed or drinking water.

**Dosage:** Spiramycin: 100 000IU/kg/day and Trimethoprim: 10 mg/kg/day  
Equivalent to 2 g of powder per 10 kg of body weight and per day during 3 to 5 days.  
The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage for spiramycin and trimethoprim per kg body weight, the amount of powder to dilute in liquid must be adjusted accordingly.

**Contraindications:** Do not use in case of known hypersensitivity to trimethoprim, spiramycin or other substance belonging to macrolide group.

**Warnings and Precautions:**

**Special warnings for each target species -** None.

**Special precautions for use**

i) **Special precautions for use in animals**

This powder for oral solution should be dissolved in milk, liquid feed or drinking water and can not be used as such.

ii) **Special precautions to be taken by the person administering the veterinary medicinal**

**product to animals**

People with known hypersensitivity to spiramycin or trimethoprim should avoid any contact with the veterinary medicinal product.

Handle the product with recommended cautions in order to avoid any risk of exposure: personal protective equipment consisting in mask, goggles, gloves should be worn when handling the product. Avoid breathing dust and avoid contact with skin and eyes.

In case of reaction after contact with product such as skin rash, seek a medical advice and show the label. Swelling of the face, of the lips or eyelids or difficulties to breath are severe symptoms and need urgent medical care.

iii) **Other precautions -** None.

**Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**Interactions with other medicaments:** None known.

**Statement on usage during pregnancy and lactation:** Laboratory studies in mice have not shown evidence of teratogenic effect for spiramycin. Foetotoxic effects have been reported in rabbits but at maternotoxic doses.

Teratogenic effects have been reported in rats at maternotoxic doses for trimethoprim, but have not been reported in rabbits.

The safety of the veterinary medicinal product has not been established during pregnancy and lactation in sows. The use of the product in pregnant or lactating sows must be submitted to a risk/benefit assessment by the responsible veterinarian.

**Adverse Effects:** None known.

**Overdose and Treatment:** None known.

**Storage Conditions:** Keep below 25°C.

**Withdrawal Period:** Meat and offal: 10 days.

Eggs: As no withdrawal period for the eggs has been established, do not use in layers when eggs are delivered to human consumption (4 weeks before the onset of the laying period and during the laying period).

**Packing:** 100g and 1kg in HDPE container.

FOR VETERINARY USE ONLY

Manufactured by :



DOPHARMA FRANCE, 23 rue du Prieuré, Saint Herblon, 44150 Vair sur Loire, France

Registration holder and repacker:



RHONE MA MALAYSIA SDN BHD (200001023279)  
Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor  
Tel: 03-7873 7355 Fax: 03-2770 0119  
E-mail: customercareline@rhonema.com

Date of revision : January 2020  
5LCRD923-100 03/XXX

W10cm H14cm

| LABEL CODE:                                     |  | VERSION                  | COLORS                                                                                                        |
|-------------------------------------------------|--|--------------------------|---------------------------------------------------------------------------------------------------------------|
| 5LCRD923-100                                    |  |                          | <div></div> Black                                                                                             |
| <b>PRODUCT:</b> CRD 92 POWDER FOR ORAL SOLUTION |  | <b>Version: 3</b>        |                                                                                                               |
| <b>PACK SIZE:</b> 100 G                         |  | <b>MMM/YY: XXX</b>       |                                                                                                               |
| <b>COUNTRY:</b> MY                              |  |                          |                                                                                                               |
| <b>DIMENSIONS:</b><br>cm = 10 x 14              |  |                          | <div>⚠ Please use the official PANTONE® (P) matching<br/>• system for an accurate color representation.</div> |
| Marketing approval:                             |  | Regulatory approval:     | GMP approval:                                                                                                 |
| <b>COMPLETE NAME:</b>                           |  | <b>COMPLETE NAME:</b>    | <b>COMPLETE NAME:</b>                                                                                         |
| <b>DATE + SIGNATURE:</b>                        |  | <b>DATE + SIGNATURE:</b> | <b>DATE + SIGNATURE:</b>                                                                                      |