

ZUPRIM 480 Water Soluble Powder

FOR ANIMAL USE ONLY

Brand or Product Name :

Zuprim 480 Water Soluble Powder

Name and Strength of Active Substance(s)

Each g contains 400 mg Sulfadiazine and 80 mg Trimethoprim.

Product Description

Zuprim 480 Water Soluble Powder is a white to off-white powder containing 40% Sulfadiazine, 8% Trimethoprim and dextrose monohydrate as diluent.

Pharmacodynamics

Mechanism of Action : Trimethoprim and sulfadiazine are antimicrobial substances which act sequentially in the synthesis of tetrahydrofolic acid from para-aminobenzoic acid. The action of trimethoprim is to potentiate the sulfadiazine component.

Pharmacodynamic properties : Veterinary preparations contain the two active substances in a 1:5 ratio and this combination is generally effective clinically because of the relatively broad range of drug ratio over which synergism occurs.

Pharmacokinetics

Pharmacokinetic properties : Both active substances are absorbed well and distributed throughout the body. Excretion occurs mainly via the urine.

Indication

Use to treat respiratory tract and alimentary tract infections caused by sulfadiazine and trimethoprim sensitive bacteria.

Recommended Dose

For administration in the drinking water. The recommended dose of Zuprim 480 Water Soluble Powder is 30mg combined actives/kg bodyweight daily, given in 2 equal doses.

Chickens: This is equivalent to 1.25g Zuprim 480 Water Soluble Powder per 20kg bodyweight birds daily. Medication period: up to 7 days.

Contraindications

None.

Warnings and Precautions

Special precautions for use

i) Special precautions for use in animals

None.

ii) Special precautions to be taken by the person administering the product to animals

Persons mixing or handling this product should avoid inhalation of any dust and contact with skin. Wear either a disposable half-mask respiratory. Rubber gloves should be worn when mixing or handling this product.

Contaminated clothing should be removed and any splashes to the skin or eyes should be washed immediately.

Hands and exposed skin should be washed thoroughly after use.

Sulphonamides may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to sulphonamides may lead to cross reaction with other antibiotics. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitive to sulphonamides.

2. If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning.

Interactions with Other Medicaments

None known.

Statement on usage during pregnancy and lactation

Not applicable.

Adverse Effects / Undesirable Effects

None recorded.

Overdose and Treatment

No treatment specified.

Withdrawal Period(s)

Poultry : 14 days.

Shelf-life

3 years from the date of manufacture. Opened bag is recommended to be used within 24 hours. Fresh medicated water must be prepared daily. Any unused medicated water must be discarded after 24 hours.

Allowable Maximum Residual Limit (MRL)

All food-producing species : Sulfa : milk (cattle) = 100 µg/kg, muscle, liver, kidney, fat (all food producing species) = 100 µg /kg. Trimethoprim : milk (cattle) = 50 µg/kg, Edible offal, muscle (chicken and mammalian) = 50 µg/kg, egg(chicken) = 50 µg/kg [Ref. REGOVP, Version 2, dated Dec 2009]

Storage Conditions

Store between 15°C - 25°C. Keep medicine out of reach of children. Jauhkan ubat dari kanak-kanak.

Disposal of Containers

Traces of medicated powder left in an opened bag can be diluted by pouring small amount of animal feed into the bag, mix well and use in animal treatment. Emptied bag can be safely disposed in trash according to local government regulations.

Dosage Forms and packaging available

Available in 100g, 200g, 500 g and 1 kg aluminum sachets.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2012-12-30