

SUFATRIM 480 Oral Suspension

FOR ANIMAL USE ONLY

Product Name :

Sufatrim 480 Oral Suspension

Name and Strength of Active Substance(s)

Each ml contains 400mg Sulfadiazine and 80mg Trimethoprim. Benzyl alcohol (as preservative) 1% w/v.

Product Description

Sufatrim 480 Oral Suspension is a white milky suspension upon agitation.

Pharmacodynamics

Alone, sulfonamides are bacteriostatic agents and trimethoprim is bactericidal, but in combination, the potentiated sulfas are bactericidal. Potentiated sulfas sequentially inhibit enzymes in the folic acid pathway, thereby inhibiting bacterial thymidine synthesis. The sulfonamide blocks the conversion of para-aminobenzoic acid (PABA) to dihydrofolic acid (DFA), and trimethoprim blocks the conversion of DFA to tetrahydrofolic acid by inhibiting dihydrofolate reductase. The trimethoprim-sulfonamides combinations have a relatively broad spectrum of activity. Gram-positive bacteria that are generally susceptible include most streptococci, many strains of staphylococci and Nocardia. Many gram-negative organisms of the family Enterobacteriaceae are also susceptible, but not *Pseudomonas aeruginosa*. Some protozoa (*Pneumocystis carinii*, Coccidia and Toxoplasma) are also inhibited by the combination. Resistance will develop slower to the combination of drugs than to either one alone.

Pharmacokinetics

Trimethoprim/sulfa is well absorbed following oral administration, with peak levels occurring about 1-4 hours after dosing. Trimethoprim/sulfa is well distributed in the body. When meninges are inflamed, the drugs enter the CSF in levels of about 50% of those found in the serum. Both drugs cross the placenta and are distributed into milk. Trimethoprim/sulfa is both renally excreted unchanged via glomerular filtration and tubular secretion and metabolized by the liver. The sulfas are primarily acetylated and conjugated with glucuronic acid and trimethoprim is metabolized to oxide and hydroxylated metabolites.

Indication

Treatment of bacterial infections due to sulfadiazine and trimethoprim sensitive organisms in poultry such as *Escherichia coli*, *Pasteurella*, *Haemophilus*, *staphylococcal* and *Salmonella* infections.

Recommended Dosage

Slowly add measured volume to water in medication tank, stirring thoroughly.

Escherichia coli, *Pasteurella*, *Haemophilus* and *staphylococcal* infections in chicken : 1ml/4.5L water.

Salmonella infections in chickens: 3ml/4.5L water.

Bacterial infections in turkey: 15ml/500kg flock liveweight in water consumed daily.

Recommended dilution rates apply when water consumption is normal; if subnormal, increase concentration to administer 15 to 30mg/kg total liveweight daily (active ingredients).

Continue treatment for 3 to 5 days according to severity of disease.

Administration

Shake the bottle well before use. Sufatrim 480 Oral Suspension disperses on dilution in water. The required dose may be added to the water header tank or to the drinking troughs.

Sufatrim 480 Oral Suspension should not be used in automatically metered water medication systems. Water systems should be clean and in good working order.

should be added to clean drinking water to provide the desired final concentration. All other sources of water should be removed during the entire period of treatment. Fresh medicated water must be prepared each day.

Contraindications

Administration to animals with known sulfonamide sensitivity, parenchymal hepatic damage or blood dyscrasia.

Warnings and Precautions

None.

Interactions with Other Medicaments

None known.

Safe Use During Pregnancy/Lactation

Do not use in laying birds producing eggs for human consumption.

Adverse Effects / Undesirable Effects

None when used as at the recommended dosage.

Overdose and Treatment

Do not exceed the stated dose.

Withdrawal Period

Poultry : slaughter 14 days. Do not use in birds producing eggs for human consumption.

Shelf-life

2 years from date of manufacture. Opened bottle is to be discarded after 24 hours. Medicated water must be prepared fresh daily. Any unused medicated water is to be discarded after 24 hours.

Allowable Maximum Residual Limit (MRL)

All food-producing species : Sulfa : milk (cattle) = 100 µg/kg, muscle, liver, kidney, fat (all food producing species) = 100 µg /kg. Trimethoprim : Edible offal, muscle (chicken and mammalian) = 50 µg/kg, egg(chicken) = 50 µg/kg [Ref. REGOV, Version 2, dated Dec 2009]

Storage Conditions

Store below 25°C. Keep medicine out of reach of children. Jauhkan ubat dari kanak-kanak.

Disposal of Container

Traces of medicated liquid left in an opened bottle can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned bottle can be safely disposed in trash according to local government regulations.

Packaging Available

Available in 1 litre per bottle, pack in 12 bottles and 20 bottles per carton.

Name and Address of Manufacturer

Life Biopharma Sdn. Bhd. (546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2013-07-10