

PACKAGE INSERT

1. PRODUCT NAME

Betamox LA 150 mg/ml Suspension for Injection

2. NAME AND STRENGTH OF ACTIVE SUBSTANCE

Active Substance:

Each ml contains:

150 mg (15% w/v) Amoxicillin

(as Amoxicillin Trihydrate 17.21 %w/v),

3. PRODUCT DESCRIPTION

Suspension for injection.

An off-white oily suspension.

4. CLINICAL PARTICULARS

4.1 Target species

Cattle, Sheep, Pigs, Dogs, Cats

4.2 Indications for use, specifying the target species

In vitro Amoxicillin is effective against a wide range of Gram-positive and Gram-negative bacteria which include:

Escherichia coli

Klebsiella pneumoniae

Proteus species

Salmonella species

Staphylococci and

Streptococci

Not effective against beta-lactamase producing organisms.

Suitable for the control of infections due to susceptible microorganisms in cattle, sheep, pigs, dogs and cats where a single injection giving prolonged activity is required. It may also protect from secondary bacterial invasion due to sensitive organisms in cases where bacteria are not the initial cause of the disease.

Indications include infections of:

- (a) Alimentary tract
- (b) Respiratory tract
- (c) Skin and soft tissue
- (d) Urogenital tract and,
- (e) In prevention of post-operative infection (treat before surgery).

4.3 Contraindications

This product is not suitable for intravenous or intrathecal use

This product should not be administered to rabbits, hamsters, gerbils or guinea pigs.

Not for use in known cases of hypersensitivity to penicillins or cephalosporins.

4.4 Special Warnings and Precautions for use

Care should be taken to avoid accidental self-injection. In the case of accidental self-injection, seek medical advice immediately. Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.
2. Handle this product with great care to avoid exposure taking all recommended precautions.
3. If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.
4. Wash hands after use.

4.5 Adverse reactions (frequency and seriousness)

Occasional local reaction may occur with use of Betamox LA.

4.6 Use during pregnancy, lactation or lay

Can be safely administered during pregnancy and lactation.

4.7 Interaction with other medicinal products and other forms of interaction

It is not generally recommended to use bactericidal and bacteriostatic antibiotics at the same time.

4.8 Amounts to be administered and administration route

To ensure the correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing.

Cattle, sheep and pigs - By intramuscular injection only.

Dogs and cats - By subcutaneous or intramuscular injection.

The recommended dosage rate is 15 mg per kg bodyweight, repeatable if necessary after 48 hours. Massage the injection site.

Shake the vial before use

Swab the septum before removing each dose. Use a dry sterile needle and syringe.

A separate injection site should be used for each administration.

Animal	Weight (kg)	Dosage volume (ml)
Cattle	450 kg	45.0 ml
Sheep	65 kg	6.5 ml
Pigs	150 kg	15.0 ml
Dogs	20 kg	2.0 ml
Cats	5 kg	0.5 ml

Dose volume is equivalent to 1 ml per 10 kg bodyweight. If dose volume exceeds 20 ml, it should be divided and injected into two sites. As with other injectable preparations normal aseptic precautions should be observed.

An appropriately graduated syringe must be used to allow accurate administration of the required dose volume. This is particularly important when injecting small volumes.

4.9 Overdose (symptoms, emergency procedures, antidotes), if necessary

Penicillins have a wide safety margin.

4.10 Withdrawal period

Milk for human consumption must not be taken from a cow during treatment. With cows milked twice daily, milk for human consumption may only be taken from 79 hours (i.e. at the 7th milking) after the last treatment. With other dosing routines, the basis of the veterinary surgeons advice should be that milk may be taken for human consumption only after the same time period from the last treatment (i.e. with three times a day milking, milk for human consumption may only be taken at the 10th milking).

Not for use in sheep producing milk for human consumption.

Animals must not be slaughtered for human consumption during treatment. Cattle may be slaughtered for human consumption only after 23 days from the last treatment. Sheep and pigs may be slaughtered for human consumption only after 16 days from the last treatment.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antibacterial

5.1 Pharmacodynamic properties

Amoxicillin predominately inhibits cell wall synthesis in susceptible bacteria. Amoxicillin has a unique mode of action which directly and irreversibly disrupts existing cell wall peptidoglycan rather than newly forming peptidoglycan of the divisory septal wall as with other members of the penicillin family.

5.2 Pharmacokinetics properties

High levels of Amoxicillin are found in kidney, urine, liver and bile after absorption due to its wide distribution. In common with other Penicillins an absence of toxicity is apparent even at high dose levels.

Animals with functional rumens should only be treated parenterally.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Butylated Hydroxyanisole
Butylated Hydroxytoluene
Aluminium Distearate
Coconut oil, Fractionated

6.2 Incompatibilities

None known

6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening the immediate packaging: 28 days

6.4 Special precautions for storage

Do not store above 25°C.

Keep out of the reach of children / Jauhi ubat dari kanak-kanak
Protect from light.

Shake the vial before use. Following withdrawal of the first dose, use the product within 28 days. Discard unused material. This product does not contain an antimicrobial preservative. Swab the septum before removing each dose.

Use a dry, sterile needle and syringe.

6.5 Nature and composition of immediate packaging

50 ml, 100 ml and 250 ml clear, colourless Type II Glass Siliconised Vials and sealed with nitril rubber bugs and aluminium caps.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MANUFACTURED BY

NORBROOK LABORATORIES LIMITED
Factory No. 4, Carnbane Industrial Estate,
Newry, Co Down, NI BT35 6QQ

8. MARKETING AUTHORISATION HOLDER

BORNEOSEW MEDICAL TRADING (M) SDN BHD
No. 57, Jalan SS 18/6, 47500 Subang Jaya
Selangor D. E., Malaysia

9. REGISTRATION NO

MALXXXXXXXXX

10. DATE OF REVISION OF THE TEXT

Date: November 2012