

CenST 240 Oral Suspension

FOR ANIMAL USE ONLY

Product Name :

CenST 240 Oral Suspension

Name and Strength of Active Substance(s) :

Each ml contains 200mg Sulfamethoxazole, 40mg Trimethoprim and 10 mg benzyl alcohol (as preservative).

Product Description :

CenST 240 Oral Suspension is a white milky suspension upon agitation.

Pharmacodynamics :

Alone, sulfamethoxazole is bacteriostatic agent and trimethoprim is bactericidal, but in combination, the potentiated sulfamethoxazole is bactericidal. Potentiated sulfamethoxazole sequentially inhibit enzymes in the folic acid pathway, thereby inhibiting bacterial thymidine synthesis. Sulfamethoxazole blocks the conversion of para-aminobenzoic acid (PABA) to dihydrofolic acid (DFA), and trimethoprim blocks the conversion of DFA to tetrahydrofolic acid by inhibiting dihydrofolate reductase. Antimicrobial activity : Both Sulfamethoxazole and trimethoprim are broad-spectrum antimicrobial agents. Their combination offer synergism against many Gram-positive and gram-negative bacteria as well as broader spectrum of activity. Gram-positive bacteria that are generally susceptible include most streptococci, many strains of staphylococci and nocardia. Many gram-negative organisms of the family Enterobacteriaceae are also susceptible. Some protozoa (pneumocystis carinii, Coccidia and Toxoplasma) are also inhibited by the combination. Resistance will develop slower to the combination of drugs than to either one alone.

Pharmacokinetics :

Trimethoprim/sulfa is well absorbed following oral administration, with peak levels occurring about 1-4 hours after dosing. Trimethoprim/sulfa is well distributed in the body. When meninges are inflamed, the drugs enter the CSF in levels of about 50% of those found in the serum. Both drugs cross the placenta and are distributed into milk. Trimethoprim/sulfa is both renally excreted unchanged via glomerular filtration and tubular secretion and metabolized by the liver. The sulfas are primarily acetylated and conjugated with glucuronic acid and trimethoprim is metabolized to oxide and hydroxylated metabolites.

Indications :

Pigs: Treatment and prevention of respiratory infections caused by Actinobacillus pleuropneumoniae susceptible to trimethoprim and sulfamethoxazole where the disease has been diagnosed in the herd.

Broilers: Treatment and prevention of respiratory infections caused by Escherichia coli susceptible to trimethoprim and sulfamethoxazole where the disease has been diagnosed in the flock

Recommended Dosage :

Must shake well before use. For oral administration via drinking water.

Pigs: 25mg combined actives per kg bodyweight per day (10.4 ml per 100 kg bodyweight per day) for 3 - 4 days.

Broilers: 33mg combined actives per kg bodyweight per day (13.75 ml per 100 kg bodyweight per day) for 3 - 4 days.

Mode of Administration :

For oral administration via drinking water.

Contraindications :

Trimethoprim/sulfa should not be used in animals showing marked liver parenchymal damage, blood dyscrasias, or in those with a history of sulfonamide sensitivity.

Warnings and Precautions :

This combination should be used with caution in animals with pre-existing hepatic disease.

Use with caution in animals with renal impairment; dosage adjustment recommended. Maintain adequate hydration to prevent crystalluria.

Interactions with Other Medicaments :

Do not combine with other veterinary medicinal products.

Safe Use During Pregnancy/Lactation :

Laboratory studies in rats conducted with trimethoprim have shown evidence of teratogenicity at higher doses than recommended therapeutic ones. The safety of the product during pregnancy and lactation has not been established in sows. The use of the product is not recommended in sows during pregnancy and lactation.

Side Effects :

A diminished water intake in chickens may occur occasionally. Hypersensitivity reactions can occur rarely.

Overdose and Treatment :

A 2 ½ fold overdose is well tolerated by pigs. In chicken an acute overdose will not occur because the birds will be reluctant to drink the strongly concentrated drinking water (too bitter taste if above 2 litres of the veterinary medicinal product per 1000 litre drinking water). Chronic overdose in chicken will result in a strongly diminished water- and feed intake and retarded growth.

Withdrawal Period :

Meat and offal: Pigs: 5 days, Broilers: 6 days

Eggs: Not authorised for use in laying birds producing eggs for human consumption.

Shelf-life :

2 years from date of manufacture. Opened bottle must be discarded after 24 hours. Medicated water must be prepared fresh daily. Any unused medicated water is to be discarded after 24 hours.

Allowable Maximum Residual Limit (MRL) :

Sulfa : milk (cattle) = 100 µg/kg, muscle, liver, kidney, fat (all food producing species) = 100 µg /kg. Trimethoprim : milk (cattle) = 50 µg/kg, Edible offal, muscle (chicken and mammalian) = 50 µg/kg, egg(chicken) = 50 µg/kg [Ref. REGOV, Version 2, dated Dec 2009]

Storage Conditions :

Store between 15°C - 25°C. Keep out of reach of children. Jauhi ubat dari kanak-kanak.

Disposal of containers :

Traces of medicated liquid left in an opened bottle can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned empty bottle can be safely disposed in trash according to local government regulations. It is recommended to send for recycling whenever possible.

Packaging :

Available in 1 litre per bottle, pack in 12 bottles or 20 bottles per carton.

Name and Address of manufacturer :

Life Biopharma Sdn. Bhd. (546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert :

2013.12.03