

CLOFENAC GEL 1% w/w

VICLOxx-03

DESCRIPTION

Colourless gel with slight alcohol odour.

COMPOSITION

Diclofenac sodium 1% w/w.

PHARMACODYNAMICS

Diclofenac Sodium is a non-steroidal anti-inflammatory drug (NSAID) with analgesic and antipyretic properties. It inhibits the activity of the enzyme cyclo-oxygenase, resulting in decreased formation of precursors of prostaglandins and thromboxanes from arachidonic acid.

PHARMACOKINETICS

Diclofenac penetrates synovial fluid, readily crosses the placenta and is excreted in breast milk. It is highly protein bound. The terminal plasma half-life is about 1 to 2 hours. It is metabolised to 4'-hydroxydiclofenac, 5-hydroxydiclofenac, 3'-hydroxydiclofenac and 4',5-dihydroxydiclofenac. It is excreted in the form of glucuronide and sulphate conjugates, mainly in the urine but also in the bile.

INDICATIONS

Non-steroidal anti-inflammatory drug indicated for the relief of pain of osteoarthritis of joints amenable to topical treatment such as knees and hands.

CONTRAINDICATIONS

Contraindicated in patients known to be hypersensitive to diclofenac sodium.

WARNINGS AND PRECAUTIONS

- Severe cutaneous reactions, including Stevens-Johnson syndrome and toxic epidermal necrolysis (Lyell's syndrome) have been reported with diclofenac sodium.
- Should only be administered onto intact skin, not on open wounds or diseased skin areas.
- Patients treated with diclofenac sodium should be closely monitored for sign of hypersensitivity reactions.
- Discontinue diclofenac sodium immediately if rash occurs.
- This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

PREGNANCY AND LACTATION

The use of diclofenac sodium is not advisable in pregnancy and lactation.

DRUG INTERACTIONS

Concomitant use of diclofenac sodium and other NSAIDs may increase the frequency of side effects.

MAIN SIDE / ADVERSE EFFECTS

- Skin rashes and eruptions have occasionally been reported and rarely urticaria.
- Cases of hair loss, bullous eruptions, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (Lyell's syndrome) and photosensitivity reactions have been reported.

OVERDOSE AND TREATMENT**Clinical features**

During recommended use there is practically no risk due to overdosage.

Treatment

Wipe off the gel and discontinue.

DOSAGE AND ADMINISTRATION

Pump Clofenac Gel 1% w/w on the affected areas 3 to 4 times per day.

Each ml of Clofenac Gel contains 9.9 mg of Diclofenac Sodium.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 30°C.

Presentation/Packing: Bottle with pump head 60 g.

Product Registration Holder /
Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman,
30010 Ipoh, Malaysia.

Revision date: November 2025

hovid