

PARA 500 g/kg Water Soluble Powder

Para 500 appears to be a white crystalline powder packed in aluminium bag with 120 micron thickness.

Composition

Each g contains

Paracetamol 500 mg

Pharmacodynamics

Paracetamol, or acetaminophen is a para amino-phenol derivative with analgesic and antipyretic properties. Its antipyretic effect may be explained by its ability to inhibit brain cyclo-oxygenases. Paracetamol is only a weak inhibitor of COX-1 synthesis and, thus, has no gastrointestinal side-effects and has no effect on platelet-aggregation.

Pharmacokinetics

Paracetamol is readily absorbed from the gastrointestinal tract. Paracetamol is metabolised mostly in the liver. The two major metabolic pathways are glucuronide conjugation and sulphate conjugation. A minor pathway catalysed by CYP (cytochrome P450) leads to the formation of the intermediary reagent, N-acetyl benzoquinoneimine which, is rapidly detoxified by reduced glutathione and removed in urine after conjugation with cystein and mercapturic acid. Paracetamol is essentially eliminated through urine (70 % of a single dose is eliminated via urine within 24 hours) under the form of paracetamol glucuronide (80 %). The other forms of elimination are cystein (10 %), unchanged paracetamol and sulphate.

Indications and Target species

In weaned pigs: Symptomatic treatment of fever in the context of respiratory diseases in combination with an appropriate anti-infective therapy, if necessary.

Dosage and administration

Administer 30 mg of paracetamol per kg of body weight per day via oral route, as long as the pigs are suffering from pyrexia, for a maximum treatment duration of 5 days. The product will be continuously administered orally in the drinking water, equivalent to 0.6 g of oral powder per 10 kg body weight. The intake of medicated drinking water depends on the clinical condition of the animals. To ensure correct dosage, the concentration in the drinking water must be adjusted accordingly. Recommendation for dissolution: Pour Para 500g/kg water soluble powder into water preferably warm (30°C to 35°C). Shake the mixture for 5 minutes to homogenize the medicated solution. Then adjust the amount of water depending on the required concentration and shake again until the solution is homogeneous.

Contraindication

Do not use in animals with known hypersensitivity to paracetamol and to any other ingredients of the product.

Do not use in animal with severe hepatic impairment.

Do not use in animal with severe renal impairment.

Do not use in animal suffering from dehydration or hypovolemia.

Warning and Precautions

Special warnings

Animals with reduced water intake and/or disturbed general condition have to be treated parenterally.

In case of combined viral and bacterial etiology of the disease, an appropriate anti-infective therapy should be given concomitantly.

Special precautions for use in animals

None

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wear appropriate protective clothing, gloves and a mask and goggles to protect the face and eyes.

If the product comes in contact with the skin or eyes, flush immediately with a large amount of water. If symptoms persist, seek medical advice.

To rule out any risk of ingestion, it is recommended not to eat, or drink while using the product and to wash the hands after use.

In the case of ingestion of the product, consult a doctor.

Do not handle the product if you are hypersensitive to paracetamol.

Interactions with Other Medicaments

Concurrent administration of nephrotoxic drugs should be avoided.

Statement on Usage During Pregnancy and Lactation

Studies in laboratory animals have not detected any teratogenic nor foetotoxic effects at therapeutic doses.

Adverse Effects /Undesirable Effects

In rare cases, at therapeutic doses, transient soft faeces can occur and can persist up to 8 days after the withdrawal of administration. It does not have any effect on general condition of animals, and resolve without any specific treatment.

Overdose and Treatment

After administration of 5 times the recommended dose of paracetamol, liquid faeces with solid particles may occasionally occur. It does not have any effect on general body condition of animals.

Acetylcysteine can be used in case of accidental overdosage.

Withdrawal Period(s)

Meat and offal: 1 day.

Storage

Store at room temperature at 30°C. Protected from light and heat exposure.

Any unused product or packaging should be disposed of in accordance with local and national legislation.

Shelf life packaged for sales: 3 years.

Shelf life after first opening: 5 days.

Shelf life after reconstitution: 24 hours.

Packaging Available

100g

Product Holder/ Manufacturer

NAM PHARM SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Date of revision

25.07.2025