

NYSTATIN FG PIL
Size: W10cm x H14cm



NYSTATIN FG Oral Powder 123.2 MIU/kg

Composition: Each kilogram contain not less than 123.2 MIU/kg Nystatin crude

Product Description: Beige colour fine powder

Pharmacodynamics: Like amphotericin B and natamycin, nystatin binds to ergosterol, a major component of the fungal cell membrane. When present in sufficient concentrations, it forms pores in the membrane that lead to K+ leakage and death of the fungus. Ergosterol is fairly unique to fungi, so the drug does not have such catastrophic effects on animals.

Pharmacokinetics: Nystatin is poorly absorbed from the GI tract, and detectable blood concentrations are not obtained after usual doses. Following oral administration, nystatin is excreted almost entirely in feces as unchanged drug.

Indication: In turkey: treatment and aid in control of crop mycosis and mycotic diarrhea (*Candida albicans*).

Routes of administration: Oral in feed.

Dosage:

For control: 50g per ton of feed.
For treatment: 100g per ton of feed.
To be fed for 7-10 days.

Contraindications: Do not use in case of hypersensitivity to Nystatin or any other component of the product.

Warnings and Precautions: People with hypersensitivity to the active ingredient or excipient should avoid contact with the product. Under normal use and handling conditions, operators should wear NIOSH approved dust mask, goggles to protect eyes and wear impermeable gloves and protective equipment to avoid direct contact with skin. Wash thoroughly with soap and water after handling. Keep out of reach of children. Drug containers and any residual contents should be disposed of safely, according to the regulation in force.

Interactions with other medicaments: None known

Statement on usage during pregnancy and lactation: The safety of the veterinary medicinal product has not been established during pregnancy and/or lactation. The use is not recommended during pregnancy and/or lactation.

Adverse Effects: None known

Overdose and Treatment: The toxicity of Nystatin is very low, even at a very high dose. There is no toxic symptoms during the administration. Nystatin does not penetrate into the cerebral or cerebrospinal fluid.

Storage Conditions: Store below 30°C. Keep away from heat and humidity

Withdrawal Period: Not required

Packing: 25 kg

Product Registration Holder:



RHONA MA MALAYSIA SDN BHD (200001023279)
Lot 18A & 18B, Jalan 241, Seksyen 51A,
46100 Petaling Jaya, Selangor.

Manufactured by:

RHONA MA MALAYSIA SDN BHD (200001023279)
Lot 16008, Jalan Nilam 3, Kawasan Perindustrian Nilai Utama,
71800 Nilai, Negeri Sembilan.
E-mail: customercareline@rhonema.com

Date of revision : XXZ
5LNYSTA3-25K 04/XX

LABEL CODE:		VERSION	COLORS
5LNYSTA3-25K			Black Please use the official PANTONE® (P) matching system for an accurate color representation.
PRODUCT: NYSTATIN FG ORAL POWDER 123.2 MIU/KG		Version: 4	
PACK SIZE: 25 KG		MMM/YY: XXX	
COUNTRY: MY			
DIMENSIONS: cm = 10 X 14			
Marketing approval:		Regulatory approval:	GMP approval:
COMPLETE NAME:		COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:		DATE + SIGNATURE:	DATE + SIGNATURE: