

SDMT 480 Water Soluble Powder

FOR ANIMAL USE ONLY

Product Name :

SDMT 480 Water Soluble Powder

Name and Strength of Active Substance(s)

Each gram of SDMT 480 water soluble powder contains Sulfadimethoxine Sodium 428 mg (which is equivalent to Sulfadimethoxine 400 mg) and Trimethoprim 80 mg.

Product Description

SDMT 480 water soluble powder is a white or almost white powder.

Pharmacodynamics

Sulfadimethoxine is a sulfonamide with long duration of action and broad spectrum of activity. It also has anticoccidial properties. It is active against Gram-positive and Gram-negative bacteria and protozoa such as coccidia. Trimethoprim belongs to the family of diaminopyrimidines. It is active against streptococci and most Gram-negative bacteria. In combination, these two active ingredients are synergistic. Sulfadimethoxine is potentiated by Diaminopyrimidine, trimethoprim. The combination of these two active ingredients enables sequential blocking of the biosynthesis of folic acid. These two substances act sequentially on the synthetic pathway of tetrahydrofolic acid: sulfonamide inhibiting the incorporation of PAB acid in folic acid, trimethoprim dehydrofolate by specifically inhibiting microbial reductase. The theoretical spectrum of activity extends to both Gram-positive bacteria (Staphylococcus, Listeria) and Gram-negative bacteria (Escherichia coli, Salmonella, Proteus, Enterobacter, Bordetella).

Pharmacokinetics

Sulfadimethoxine is considered acting sulfonamide with a longer persistence of plasma levels. Its binding to plasma proteins is important. The distribution is good in most tissues and organs. Trimethoprim is rapidly absorbed after oral administration. It is widely distributed in the body. The two active ingredients are partially metabolized in the liver. Their elimination is primarily renal.

Indication

Infections sensitive to trimethoprim and sulfadimethoxine.

Calves, pigs and poultry :

- Treatment and prevention of respiratory and gastrointestinal infections.
- Treatment and prevention of digestive coccidiosis.

Recommended Dosage

- 18.68 to 37.36 mg of sulfadimethoxine and 4-8 mg of trimethoprim per kg of body weight per day or

- 0.05 g to 0.1 g **SDMT 480 Water Soluble Powder** per kg of body weight per day

Use for 3 to 5 consecutive days. Adjust according to actual consumption of animals to meet the weight dose (mg/kg).

Mode of Administration

To be administered via animal drinking water. Prepare fresh medicated solution daily.

Contraindications

Do not use if you are allergic to sulfonamides or trimethoprim. Do not use in animals with severe renal or hepatic insufficiency. In the absence of a waiting time for the eggs do not use in laying birds producing eggs for human consumption (4 weeks before the start of lay and during it).

Warnings and Precautions

Precaution for animals :

Not for use in animals who are hypersensitive (allergic) to it or other sulfa medications. Not for use in pregnant or lactating animals (female animals nursing their young). The animals should be encouraged to drink water to prevent crystals from forming in the urine.

Precaution for person handling the product :

Use appropriate personal protection. Wear impervious gloves when handling powder. Open sachet with care and not in windy area to avoid forming dust cloud. Use adequate masks and safety goggles in case of powder being blown.

Respiratory Protection	:	Wear dust mask in case powder being blown.
Hands	:	Wear Impervious gloves when handling powder.
Eye Protection	:	Wear Safety goggles when handling powder.
Skin Protection	:	Wear protective clothing to minimize skin exposure.

Interactions with Other Medicaments

Unknown.

Pregnancy and Lactation

Teratogenic and fetotoxic effects have been observed in laboratory animals at higher than recommended therapeutic doses. Use of the product during pregnancy and lactation is inappropriate.

Adverse Effects

Side effects due to potentiated sulfonamides are quite rare, and adverse reactions can mainly attributed to the sulfonamide. The more common adverse effects are urinary tract disturbances (crystalluria, hematuria, or even obstruction), hematopoietic disorders (thrombocytopenia, anemia, leucopenia) and dermatologic reaction.

Overdose and Treatment

None known.

Withdrawal Period(s)

Meat and offal: 12 days. In the absence of a waiting time for the eggs do not use in laying birds producing eggs for human consumption (4 weeks before the start of lay and during it).

Shelf-life

Shelf life as packaged for sale: 3 years

Shelf life after first opening of the immediate packaging: 24 hours

Shelf life after reconstitution in drinking water: 24 hours

Allowable Maximum Residual Limit (MRL)

Sulfadimethoxine : Cattle (fat, kidney, liver, muscle), Chickens (kidney, liver, muscle, skin and fat), Swine (kidney, liver, muscle, skin and fat) = 100 µg /kg, Milk of cattle : 10µg /kg.

[Ref. Health Canada. May 22 2013]

Trimethoprim : Milk (cattle), Edible offal, muscle (mammalian and chicken), egg (chicken) = 50 µg /kg

[Ref. REGOVP, Version 2, dated Dec 2009]

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhi ubat daripada kanak-kanak.

Disposal of containers

Empty aluminum sachet must be safely disposed according to local government regulations.

Dosage Forms and packaging available

Available in 200g, 500g 1000g powder per sachet, pack in 10, 20, 50 and 100 sachets per box.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2014-04-14