

Nystatin 50g/kg Premix Powder

FOR ANIMAL USE ONLY

Brand or Product Name :

Nystatin 50g/kg Premix Powder

Name and Strength of Active Substance(s)

Each kilogram contains 50 g of Nystatin (minimum 100,000 IU/g).

Product Description

Nystatin 50g/kg Premix Powder is a light brownish powder to be used in animal feed.

Pharmacodynamics

Nystatin is polyene antifungal antibiotic produced by *Streptomyces noursei*. Nystatin acts by binding to sterols in the membrane of the fungal cell where it alters the permeability of the fungal cell membrane allowing intracellular potassium and other cellular constituents to leak out.

Nystatin has activity against a variety of fungal organisms, many molds and yeast, including *Candida* spp.

Pharmacokinetics

Nystatin is not measurably absorbed after oral administration and is almost entirely excreted unchanged in feces.

Indications

For the control and treatment of crop mycosis and mycotic diarrhea (*Candida albicans*) in turkeys.

Route of Administration

Per oral via animal feed.

Recommended Dosage

For growing turkeys :

Aid in control of crop mycosis and mycotic diarrhea : 1 kg (equivalent to 50g Nystatin) per ton of feed

For treatment of crop mycosis and mycotic diarrhea : 2 kg (equivalent to 100g Nystatin) per ton of feed, to be fed for 7 to 10 days.

Direction of Use

To ensure thorough dispersion of the product it should first be mixed with a suitable quantity of feed before incorporation into the final mix.

Contraindication

Do not use in known cases of hypersensitivity to active ingredient.

Warning and Precautions

Use under veterinarian recommendation.

Drug Interactions

No significant interactions reported for oral nystatin.

Pregnancy and lactation

The lack of appreciable oral absorption of nystatin makes it safe to use during pregnancy and lactation.

Adverse Effects

Occasionally, nystatin may cause gastrointestinal upset (anorexia, vomiting, diarrhea) when administered in high dosages.

Overdose and Treatment

Because the drug is not absorbed after oral administration, acute toxicity after an oral overdose is extremely unlikely, but transient gastrointestinal distress may result.

Withdrawal Period(s)

0 day.

Shelf-life

Original Packing : 2 years from the date of production.

Opened bag : 24 hours.

Reconstituted feed : recommended to be used within 24 hours.

Maximum Residual Limit (MRL)

Edible tissue (pig and poultry), egg (poultry) = 0mg/kg [Ref. REGOVP, Version 2, dated Dec. 2009]

Storage Conditions

Store in a dry place below 25°C. Keep medicine out of reach of children. Jauhkan ubat daripada kanak-kanak.

Disposal of Containers

Emptied bag must be safely disposed according to local government regulations.

Dosage Forms And Packaging Available

Available in 25 kg PP laminated Paper bags.

Name and Address of Manufacturer

Life Biopharma Sdn. Bhd. (Company No. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan Darul Khusus, Malaysia.

Date of Revision of Package Insert

2014-03-14