

ERYCIN 300 g/kg Water Soluble Powder

Erycin 300 appears to be a pink crystalline powder packed in aluminum bag with 120 micron thickness.

Composition

Each kg contains
Erythromycin Thiocynate
300 g

Pharmacodynamics

Macrolides are bacteriostatic, with potential for a time-dependent bactericidal action, particularly with high concentrations. Macrolides are thought to enter the cell and reversibly bind to the 50 S ribosomal subunit, inhibiting translocation of peptides, thereby inhibiting protein synthesis. Although macrolides bind to mitochondrial ribosomes, like chloramphenicol, macrolides are unable to cross the mitochondrial membrane and so do not produce bone marrow suppression in mammals.

Pharmacokinetics

Erythromycin is absorbed rapidly when administered orally and diffusion occurs into most tissues and fluids. Elimination of erythromycin occurs primarily through hepatic metabolism and the remainder is excreted in active form in the urine and bile.

Indications

For the treatment of erythromycin susceptible infections of poultry, including the Chronic Respiratory Disease Complex (CRD), Coryza (*Haemophilus paragallinarum*), and Cholera (*Pasteurellosis*).

Directions for Use

30mg per kilogram bodyweight daily.

Contraindication:

Do not use in known cases of hypersensitivity to active ingredient.

Mode of Administration

Oral Water soluble powder.

Warning and Precautions

Use under veterinarian recommendation.

Interactions with Other Medicaments

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other products.

Statement on Usage During Pregnancy and Lactation

None known.

Adverse Effects / Undesirable Effects

If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

Overdose and Treatment

Stop medication and to seek for veterinarian advice if there is sign of over dosage.

Withdrawal Period (s)

Chicken: 6 days

Eggs: 6 days

Storage Condition

Store below 30 degrees Celsius. Protect from heat and sunlight exposure.

Keep out of reach of children (Jauhkan dari jangkauan kanak-kanak)

Shelf-life after opening: 7 days

Shelf-life after reconstitution: 24 hours

Any unused veterinary product or waste materials derived from such veterinary medicinal products should be disposed of an accordance with local requirements

Date of Revision

06-08-2019

Packaging Available

100g & 160 g.

Name and Address of Manufacturer

NAM PHARM SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833