

NAMCILLIN-50% w/w Water Soluble Powder

Namcillin-50 appears to be white or almost white crystalline powder.

Composition:

Each 100g contains
Amoxicillin Trihydrate 57.5 g

Pharmacodynamics:

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against gram positive and gram negative bacteria. It owes its activity to the inhibition of the development of peptidoglycan network structure in the bacterial cell wall.

Pharmacokinetics:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Indications:

Chickens: For the treatment of pasteurellosis and colibacillosis.

Turkeys: For the treatment of pasteurellosis.

Ducks: For the treatment of infections caused by Streptococcus bovis, Pasteurella anatipestifer and Escherichia coli.

Pigs: For the treatment of salmonellosis and pasteurellosis caused by isolates sensitive to amoxicillin.

Directions for Use:

The product is administered in the drinking water. Prepare the solution with fresh tap water immediately before use. Once opened, use the contents of one sachet immediately. In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

Chickens

The recommended dosage is 15 mg amoxicillin trihydrate per kg bodyweight.

The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks

Recommended dosage is 20 mg amoxicillin trihydrate/kg bodyweight for 3 consecutive days.

Turkeys

Recommended dosage is 15-20 mg amoxicillin trihydrate/kg bodyweight for 3 days or in severe cases for 5 days.

Swine

Administer in the drinking water to give 20mg/kg bodyweight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days.

Mode of Administration:

Oral

Contraindication:

Namcillin-50 should not be administered to rabbits, hamsters, gerbils and guinea pigs. Do not use in animals with known hypersensitivity to the active substance.

Warning and Precautions:

(i) Special precautions for use in animals

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Avoid inhalation of dust.

Wash hands after use.

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1) Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.

2) Handle this product with great care to avoid exposure, taking all recommended precautions.

3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments:

None known.

Statement on Usage During Pregnancy and Lactation:

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

Use only according to the benefit/risk assessment of the responsible veterinarian.

Adverse Effects / Undesirable Effects:

None known.

Overdose and Treatment:

There are no updated known effects at overdose. Stop medication and to seek for veterinarian advice if there is sign of overdosage.

Withdrawal Period (s):

Meat:

Chickens 1 day

Ducks 9 days

Turkeys 5 days

Pigs 2 days

Not authorised for use in laying birds producing eggs for human consumption

Packaging Available: 100 g.

Storage Condition:

Store below 30 degrees Celsius. Protect from heat and sunlight exposure. Keep out of reach of children (Jauhkan dari kanak-kanak).

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf-life after opening: 6 days

Shelf life after reconstitution: 24 hours

Date of revision: 31-03-2014

Name and Address of Manufacturer:

NAM PHARM SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833