

ALBAZOL 10% Oral Suspension

FOR ANIMAL USE ONLY

Brand or Product Name :

Albazol 10% Oral Suspension

Name and Strength of Active Substance(s)

Each ml contains Albendazole 100mg, Methylparaben 0.3% as preservative, Propylparaben 0.03% as preservative

Product Description

Albazol 10% Oral Suspension is a milky white suspension after shaking well.

Pharmacodynamics

A broad spectrum multi-purpose anthelmintic for the control of mature and developing immature forms of gastrointestinal roundworms, lungworms, tapeworms and adult liver fluke in cattle and sheep. The product is also ovicidal against fluke and roundworm eggs.

Pharmacokinetics

Benzimidazoles bind to nematode tubulin, a protein necessary for the formation and viability of microtubules. This occurs primarily in absorptive intestinal cells resulting in the absence of microtubules in the intestinal cells of the nematode, with the result that these cells cannot absorb nutrients, thus causing a consequent reduction in glycogen and effective starvation of the parasites. Structural differences have been shown to exist between tubulin from mammalian and helminth sources, resulting in the preferential toxicity of albendazole to the helminth and not to the host. Benzimidazoles have also been shown to inhibit the fumarate reductase system of helminths and impair energy production.

Indication

A broad spectrum multi-purpose anthelmintic for the control of mature and developing immature forms of gastrointestinal roundworms, lungworms, tapeworms and adult liver fluke in cattle and sheep. The product is also ovicidal against fluke and roundworm eggs.

In cattle it is active against the following species: Roundworms: Ostertagia, Haemonchus, Trichostrongylus, Nematodirus, Oesophagostomum, Bunostomum, Cooperia, and Strongyloides spp. It is usually effective against inhibited larvae of Cooperia and Ostertagia, Lungworms: Dictyocaulus viviparus, Tapeworms: Moniezia spp., Adult liver fluke: Fasciola hepatica. In sheep it is active against benzimidazole-susceptible strains of the following species: Roundworms: Ostertagia, Haemonchus, Trichostrongylus, Nematodirus (including N. battus), Chabertia and Oesophagostomum. It is usually effective against inhibited larvae of Ostertagia, Lungworms: Dictyocaulus filaria, Tapeworms: Moniezia spp., Adult liver fluke: Fasciola hepatica. The product is ovicidal and will kill fluke and roundworm eggs, thus reducing pasture contamination.

Recommended Dosage

Shake well before use. For oral administration only using properly calibrated dosing equipment. To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked. One ml of the product contains 100mg Albendazole.

Cattle : Worm dose : For the control of roundworms, lungworms, tapeworms and fluke and roundworm eggs. Dosage : 7.5mg albendazole per kg bodyweight.

Fluke and worm dose : For the additional treatment of adult liver fluke (chronic fascioliasis) in cattle. Dosage : 10mg albendazole per kg bodyweight.

Sheep : Worm dose : For the control of roundworms, lungworms, tapeworms, fluke and roundworm eggs. Dosage : 5 mg albendazole per kg bodyweight.

Fluke and worm dose : For the additional treatment of adult liver fluke (chronic fascioliasis) in sheep. Dosage : 7.5 mg albendazole per kg bodyweight.

Mode of Administration

Shake well before use. For oral administration only using properly calibrated dosing equipment.

Contraindications

Known hypersensitivity to the active ingredient.

Warnings and Precautions

Special warnings for each target species

Cattle suffering from severe lung damage due to heavy lungworm infestation may continue to cough for some weeks after treatment. Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.

- Underdosing, which may be due to underestimation of bodyweight, misadministration of the product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used. Resistance to benzimidazoles (which includes albendazole) has been reported in Teladorsagia, Haemonchus, Cooperia and Trichostrongylus species in small ruminants in a number of countries including the EU. Resistance to albendazole has been reported in Cooperia and Teladorsagia species in cattle in developed countries such as New Zealand. Therefore the use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics

Special precautions for use

i. Special precautions for use in animals : Not to be diluted or mixed with other products. Avoid the introduction of contamination during use. Care must be taken not to damage the pharyngeal region during dosing, particularly in sheep. Intensive use or misuse of anthelmintics can give rise to resistance. To reduce the risk, dosing programmes should be discussed with a veterinary surgeon.

ii. Special precautions for the person administering the veterinary medicinal product to animals : Direct contact with the skin should be kept to a minimum. Wear suitable protective clothing including impermeable rubber gloves. In the event of accidental eye exposure, flush eye thoroughly with running water. If irritation persists, seek medical attention. In the event of accidental skin exposure, wash the affected area with soap and water. If irritation persists, seek medical advice. Wash hands after use.

iii. Other precautions : None

Interactions with Other Medicaments

Unknown.

Pregnancy and lactation

Do not dose ewes at the 'fluke and worm' dose rate, (7.5 mg/kg), during tupping or for 1 month after removing the rams. Can be safely used during lactation. The use of the product in breeding bulls or pregnant cattle is not expected to interfere with their reproductive performance.

Adverse Effects

None known.

Symptoms and treatment of Overdose

None known. Do not exceed recommended dose.

Withdrawal Period(s)

Cattle: Meat & Offal: 14 days. Milk: 60 hours. Sheep: Meat & Offal: 5 days. Not to be used in sheep producing milk for human consumption.

Shelf-life

2 years from date of manufacture. Opened bottle must be discarded after 24 hours. Medicated water must be prepared fresh daily. Any unused medicated water is to be discarded after 24 hours.

Allowable Maximum Residual Limit (MRL)

Muscle, fat (cattle and other species), milk (cattle) = 100 µg/kg ; Liver, kidney (cattle and other species) = 5000 µg/kg [Ref. REGOV, Version 2-Dec 2009]

Storage Conditions

Store airtight, away from sunlight and below 30°C. Keep out of reach of children. Jauhi ubat daripada kanak-kanak.

Disposal of containers :

Traces of medicated liquid left in an opened bottle can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned empty bottle can be safely disposed in trash according to local government regulations. It is recommended to send for recycling whenever possible.

Packaging

Available in 1 litre per bottle, pack in 12 bottles or 20 bottles per carton.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2014-04-02