

Package Insert**Brand or Product Name**

OXITETRACICLINA 200mg/ml L.A. Solution for Injection
Oxytetracycline

Name and Strength of Active Substance(s)

Oxytetracycline Dihydrate 215.6mg/ml

Product Description

A transparent, yellow, slightly dense and viscous solution.

Pharmacodynamics

Oxytetracycline is a bacteriostatic antibiotic that inhibits protein synthesis in susceptible bacteria. Inside the cell it binds irreversibly to receptors on the 30S subunit of the bacterial ribosome where it interferes with the binding of the aminoacyl-transfer RNA to the acceptor site on the messenger RNA ribosome complex. This effectively prevents the addition of amino acids to the elongating peptide chain, inhibiting protein synthesis. The product is specifically formulated to provide a prolonged action resulting in sustained antibacterial activity. Oxytetracycline had been shown to be effective in vitro against the following bacterial species: Bordetella bronchiseptica, Corynebacterium pyogenes, Erysipelothrix rhusiopathiae, Escherichia coli, Haemophilus somnus, Pasteurella haemolytica, Pasteurella multocida, Salmonella dublin, Staphylococcus aureus, Streptococcus agalactiae, Streptococcus faecalis, Streptococcus pyogenes and Streptococcus uberis.

Pharmacokinetics

Blood levels persist for at least 4 days after administration by the intramuscular route. Maximum blood levels are achieved between 4 and 8 hours following intramuscular administration.

Environmental Properties

not relevant

Indication

The product is indicated for use in cattle, sheep and pigs in the treatment of: - Atrophic rhinitis caused by Bordetella bronchiseptica, Mannheimia haemolytica and Pasteurella multocida. - Navel/joint ill caused by Arcanobacterium pyogenes, E. coli or Staphylococcus aureus. - Mastitis caused by Corynebacterium pyogenes, E. coli, Staphylococcus aureus, Streptococcus agalactiae or Streptococcus uberis. - Metritis caused by E. coli or Streptococcus pyogenes. - Pasteurellosis and infections of the respiratory tract caused by Mannheimia haemolytica and Pasteurella multocida. - Septicaemia caused by Salmonella dublin and Streptococcus pyogenes. - Erysipelas caused by Erysipelothrix rhusiopathiae. The product can also be used in the control of enzootic abortion in sheep.

Recommended Dosage and Mode of Administration**Intramuscular use.**

The recommended dose rate is 20 mg/kg bodyweight (i.e. 1 ml per 10 kg bodyweight) administered by deep intramuscular injection. The product is recommended for a single administration only. Maximum recommended dose at any one site: Cattle : 20ml Pigs : 10ml Sheep : 5ml Piglets : 1 day 0.2ml 7 days 0.3ml 14 days 0.4ml 21 days 0.5ml over 21 days 1.0 ml/10kg.

Mode of Administration

Intramuscular route

Contraindications

Not for use in horses, dogs, cats. Not for use in animals suffering from hepatic or renal damage.

Warnings and Precautions

No special warnings. Special Precautions for use in animals: Do not dilute the product. If concurrent treatment is administered, use a separate injection site. Special precautions to be taken by the person administering the product to the animals: Wash hands after use. In case of contact with eyes or skin, wash immediately with plenty of water as irritation may occur. Take care to avoid accidental injection

Interactions with Other Medicaments

None known.

Pregnancy and lactation

The use of oxytetracycline during the period of tooth and bone development, including late pregnancy may lead to tooth discoloration, the product can be safely administered to lactating animals.

Adverse Effects/Undesirable Effects

Although the product is well tolerated, occasionally a slight local reaction of a transient nature has been observed.

Overdose and Treatment

There is no known specific antidote, if signs of possible overdose occur treat the animal symptomatically.

Withdrawal Period(s)

Cattle: Meat – 31 days Milk – 10 days Sheep: Meat – 9 days Milk – 7 days Pigs: Meat – 18 days

Storage Conditions

Do not store above 25°C. Protect from light. Eventually, the preparation may get dark, which does not imply loss of activity.

Shelf Life After First Opening

28 days.

Incompatibilities

None known.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

Dosage Forms

Solution for injection.

Packaging Available

Vials of 100ml

Name and Address of Manufacturer

Industrial Veterinaria, S.A.
Esmeralda,19
E-08950 Esplugues de Llobregat (Barcelona) Spain.

Marketing Authorisation Holder

Taseen Trading Sdn. Bhd.
No 8, Jalan Astana 1F/KU2,
Bandar Bukit Raja, 41050,
Klang, Selangor DE. Malaysia

Date of Revision of Package Insert

November 12, 2018-April 12, 2019