

INVEMOX 15% W/V LA

Long-Acting injectable suspension Amoxicillin

Name and strength of active substance

Amoxicillin (as trihydrate), 150 mg/ml.

Indications

The product is suitable for the control of infections, due to susceptible micro-organisms, in cattle, pigs, sheep, dogs and cats where a single injection giving prolonged antibiotic cover is required. It may also protect from secondary bacterial invasion in cases where bacteria are not the initial cause of the disease.

Indications include infections of: Alimentary tract, respiratory tract, skin and soft tissue, urogenital tract and in prevention of post-operative infection (treat before surgery). In vitro amoxicillin is effective against a wide range of Gram-positive and Gram-negative bacteria which include: *Escherichia coli*, *Klebsiella pneumonia*, *Proteus spp.*, *Salmonella spp.*, *Staphylococci* and *Streptococci*.

Contraindications

This product is not suitable for administration via intravenous or intrathecal routes. Not to be administered to animals sensitive to penicillin. Not to be used in rabbits, guinea pigs, hamsters, gerbils or in any other small herbivores. Not effective against beta-lactamase producing organisms.

Adverse effects

Occasional local tissue reaction may result from use of this product.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

Target species

Cattle, sheep, pigs, dogs and cats.

Recommended dosage and mode of administration

Cattle, sheep and pigs: Intramuscular injection only.

Dogs and cats: Intramuscular or subcutaneous injection. 1 ml of INVEMOX 15% LA per 10 kg b.w. (equivalent to 15 mg of amoxicillin/kg of b.w.).

Repeated if necessary after 48 hours. A separate injection site should be used for each administration.

Suggested doses are: Cattle 500 kg - 50 ml; Sheep 50 kg - 5 ml; Pigs 50 kg - 5 ml; Dogs 10 kg - 1 ml; Cats 5 kg - 0.5 ml. An appropriately graduated syringe must be used to ensure accurate administration of the required dose volume. This is particularly important when injecting small volumes.

Shake the vial before use. Use a dry, sterile needle and syringe. After administration massage the injection site.

Dose volumes in excess of 20 ml should be injected at more than one site.

Withdrawal Period

Meat: Animals must not be slaughtered for human consumption during treatment. Cattle may be slaughtered for human consumption only after 23 days from the last treatment.

Sheep and pigs may be slaughtered for human consumption only after 16 days from the last treatment.

Milk: Milk for human consumption must not be taken from a cow during treatment. With cows milked twice daily, milk for human consumption may only be taken from 79 hours (i.e. at the 7th milking) after the last treatment. With other dosing routines, the basis of the veterinary surgeon's advice should be that milk may be taken for human consumption only after the same time period from the last treatment (i.e. with three times a day milking, milk for human consumption may only be taken at the 10th milking). Not for use in sheep producing milk for human consumption.

Storage conditions

Keep out of the sight and reach of children.

Store below 30 °C.

Shelf life after first opening: 28 days.

Special warnings

Special warning for each target species: Not to be used in animals with previous sensitivity to penicillins.

Special precautions for use in animals: None.

Special precautions to be taken by the person administering the medicinal product to the animals: Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may cause cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Handle this product with great care to avoid exposure, taking all recommended precautions. If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention. Wash hands after use.

Statement on usage during pregnancy and lactation: No special precautions are necessary.

Interactions with other medicaments: Amoxicillin exerts its

bactericidal action by inhibition of bacterial cell wall synthesis during multiplication. It is therefore in principle not compatible with bacteriostatic antibiotics (tetracyclines, chloramphenicol) which inhibit multiplication. Synergism occurs with β -lactam antibiotics and aminoglycosides.

Overdose and treatment: Amoxicillin has a wide safety margin. In case of severe allergic reactions, treatment should be discontinued and corticoids and adrenaline should be administered. In the rest of cases, symptomatic treatment should be administered.

Incompatibilities: None known.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

Other information

Product description: A creamy-white oily suspension. Homogeneous at shake, it is easily redispersed.

Pharmacodynamics: The active ingredient, amoxicillin, is a bactericidal antibiotic of the β -lactam class which acts by inhibition of bacterial cell wall synthesis. Amoxicillin is not resistant to the action of beta-lactamases which can hydrolyse the molecules causing the beta-lactam ring structure to open, rendering it antibiotically inactive. Due to its wide distribution after absorption, high levels of amoxicillin are found in kidney, urine, liver and bile.

Pharmacokinetics: Following intramuscular injection in cattle at the recommended dosage rate peak levels were observed 2.98 hours following administration. Following intramuscular injection in sheep at the recommended dosage rate peak levels were observed 0.96 hours following administration. Following intramuscular injection in pigs at the recommended dosage rate peak levels were observed 1.18 hours following administration.

Packaging available: 100 ml.

For animal treatment only. To be supplied only on veterinary prescription.

Name and address of manufacturer

Industrial Veterinaria, S.A.
Esmeralda, 19
08950 Esplugues de Llobregat
(Barcelona) Spain.

Marketing authorization holder

Taseen Trading Sdn. Bhd.
No 8, Jalan Astana 1F/KU2
Bandar Bukit Raja, 41050,
Klang, Selangor DE. Malaysia

Date of revision of package insert
April 12, 2019