

SULPASTA WATER SOLUBLE POWDER

Sulpasta water soluble powder appears to be a white or almost white crystalline powder.

Composition

Each 100g contains

Sulfadiazine sodium	40 g
Trimethoprim	8 g

Pharmacodynamics

Veterinary preparations contain the two active substances in a 1:5 ratio and this combination is generally effective clinically because of the relatively broad range of drug ratio over which synergism occurs.

Pharmacokinetics

Both active substances are absorbed well and distributed throughout the body. Excretion occurs mainly via the urine.

Indications

Swine: For the treatment and control of diseases sensitive to trimethoprim/sulfadiazine in pigs, in particular the respiratory disease complex.

Chickens: For the treatment and control of diseases sensitive to trimethoprim/sulfadiazine in chickens, in particular enteric and respiratory bacterial infections.

Dosage and administration

Oral administration.

For administration in the drinking water, the recommended dose of Sulpasta is 30 mg combined actives/kg bodyweight daily, given in 2 equal doses.

Swine: This is equivalent to 40 g Sulpasta per 640 kg pig weight daily. Medication period: 5 days.

Chickens: This is equivalent to 1 g Sulpasta per 16 kg bodyweight birds daily. Medication period: up to 7 days.

The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of the active substances may have to be adjusted accordingly. To allow ready dissolution of the Sulpasta, dissolve the required amount

of product in a few liters of water before addition to the header tank. Avoid any further dilution by securing the header tank ballcock until all the medicated water has been consumed. Once medicated water has been consumed, the ballcock can be released and non-medicated water re-supplied. Any medicated water which is not consumed within 24 hours should be discarded.

Contraindication

Hypersensitivity to trimethoprim and/or sulphonamides. Administration to animals with a serious impaired renal and/or liver function or with blood dyscrasias.

Warning and Precautions

None known.

Interactions with Other Medicaments

None known.

Statement on usage during pregnancy and lactation

In the absence of toxicology study, this veterinary medicinal product must not be used during pregnancy and lactation. Do not use in laying birds producing eggs for human consumption.

Adverse Effects

If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

Overdose and Treatment

Stop medication and to seek for veterinarian advice if there is sign of overdosage.

Withdrawal Period (s)

Animals and birds must not be slaughtered for human consumption during treatment.

Swine: Pigs may be slaughtered for human consumption only after 14 days from the last treatment.

Chickens: Chickens may be slaughtered for human consumption only after 7 days from the last treatment.

Eggs from treated chickens must not be used for human consumption.

Storage

Store below 30°C. Protect from heat and sunlight exposure.

Keep medicine out of reach of children (Jauhkan ubat daripada kanak-kanak).

Any unused product or packaging should be disposed of in accordance with local legislation.

Shelf-life after opening: 6 days.

Shelf-life after reconstitution: 24 hours.

Packaging

100 g

Manufacturer and product holder

NAM PHARM SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Date of revision

23-02-2026