

Product Name:

SULPHADIAZINE & TMP Water Soluble powder

Composition:

Each gram contains:

Sulphadiazine	400mg
Trimethoprim	80mg

Product Description:

Light Yellow powder

Pharmacodynamics:

Sulfadiazine is a bacterial antibiotic belonging to the sulphonamide group, which acts by interference with the synthesis of nucleic acids. Trimethoprim is a dihydrofolate reductase inhibitor which also interferes with the synthesis of bacterial nucleic acids. Sulfadiazine and trimethoprim act on the same metabolic pathway, resulting in potentiation of antibacterial activity.

Pharmacokinetics:

Following oral administration of sulfadiazine and trimethoprim to chickens, t_{α} and t_{β} values of 0.756 and 7.07 hours (sulfadiazine) and 0.680 and 6.24 hours (trimethoprim) were obtained. Values for t_{max} were 2.46 and 2.44 hours, values for C_{max} were 86.45 and 3.65g/ml and values of AUC were 620.50 and 19.87g hour/ml, respectively for sulfadiazine and trimethoprim.

Both active substances are absorbed well and distributed throughout the body. Excretion occurs mainly via the urine.

Indications:

Bacterial infections due to susceptible organisms in chickens, turkeys and pigs.

Dosage and Administration:

Poultry: For oral administration in drinking water. E.coli, Pasteurella, Actinobacillus and Staphylococcal infections: Add 100-300g/350L of drinking water for 3-5 days. Pigs: For oral administration in drinking water. For enteric infections and pneumonia: Add 100-300g/350L of drinking water for 3-5 days.

Contraindications:

Not to be used in animals with known sensitivity to the active ingredients.

Interaction with Other Medicaments:

None known.

Pregnancy and Lactation:

In the absence of toxicology studies, this veterinary medicinal product must not be used during pregnancy and lactation. Do not use in laying birds producing eggs for human consumption.

Side effects:

Hypersensitivity to sulphonamides has been rarely reported, however this remains a potential undesirable effect. Crystalluria caused by precipitation of insoluble sulphonamide is a theoretical risk although rarely reported.

Symptoms and Treatment of Overdose:

No information available. As there is no specific antidote, treatment should be symptomatic.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Use under veterinarian recommendation.

Withdrawal Periods:

Poultry and Pigs: Do not administer later than 14 days before slaughter for human consumption.

Do not use in laying birds where eggs or egg products are to be used for human consumption or processing.

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,

Prai Industrial Estate MK1,

13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahan Klebang II,

Taman Perindustrian Antarabangsa IGB

31200 Chemor, Perak.

DATE OF REVISION: APRIL 2019