

Hampiseptol Solution for Injection

Hampiseptol Solution for Injection appears to be a clear and colourless solution

Composition:

Each 1 mL contains	
Sulfadimidine sodium	200 mg
Trimethoprim	40 mg

Pharmacodynamics:

Sulfamethazine is a sulfonamide fast action in the body. It has only a bacteriostatic activity, blocking the use of folic acid by the bacteria. It has a broad spectrum of activity ranging from Gram-positive to Gram-negative organisms (streptococci, staphylococci, corynobactéries, Pasteurella and E. coli.).

Trimethoprim belongs to the family of diaminopyrimidines. It is active against streptococci and most Gram-negative bacteria, trimethoprim specifically inhibits microbial reductase dehydrofolate. In combination these two active substances are synergistic. The association allows a sequential blocking of biosynthesis of tetrahydrofolic acid, sulfonamide inhibiting incorporation of para-aminobenzoic acid in the folic acid, trimethoprim by specifically inhibiting microbial dehydrofolate reductase acid. The theoretical activity spectrum and extends to both Gram positive (Staphylococcus, Listeria.) And Gram negative (Escherichia coli, Salmonella, Proteus, Enterobacter, Bordetella).

Pharmacokinetics:

The SMZ is considered a semi-sulfonamide delay with enough long persistence of plasma levels. Its binding to plasma proteins is important. The distribution is good in most tissues and organs. Its elimination is mainly urinary. Trimethoprim is rapidly absorbed after oral administration. It is widely distributed in the body.

Indications:

Systemic treatment for a range of bacterial infections caused by bacterial sensitive to trimethoprim + sulfadimidine of the respiratory tract (pneumonia, bronchitis), urogenital tract (nephritis, metritis), gastrointestinal tract (colibacillosis, salmonellosis).

Directions for Use:

Horses: 10mL per 200kg bw daily by intravenous (IV) injection only.

Cattle, Sheep & Pigs: 1mL per 10-15 kg bw daily by intramuscular (IM) or intravenous (IV) injection.

Dogs: 1mL per 8kg bw daily by intramuscular (IM) injection.

Cats: 1/2mL by intramuscular injection daily.

In severe cases dosage should be repeated for 1-3 days.

Transitory pain may be experienced following large volume injection.

Doses of more than 10mL should be injected at separate sites.

Mode of Administration:

intramuscular (IM) or intravenous (IV)

Incompatibilities:

None known

Contraindication:

Do not administer to animals hypersensitive to sulfonamides and / or trimethoprim.

Do not use in patients with renal or hepatic insufficiency serious.

Warning and Precautions:

Special warnings for each target species

No.

Special precautions for use in animals

Divide the doses in several points during injection intramuscular administrations.

Special precautions for the person administering the veterinary medicinal product to animals

People with known hypersensitivity to sulfonamides or trimethoprim should avoid contact with the veterinary medicinal product.

Accidental contact with eyes, rinse thoroughly with water.

Consult a doctor if skin rash.

Consult a doctor immediately in case of edema of the face, lips or eyes or difficulty breathing occur.

Interactions with Other Medicaments:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other products.

Statement on Usage During Pregnancy and Lactation:

Teratogenic and fetotoxic effects have been observed in laboratory animals at higher than recommended therapeutic doses. Use of the product during pregnancy is contraindicated.

Adverse Effects / Undesirable Effects:

Local reactions at the injection site after intramuscular administration.

Overdose and Treatment:

Stop medication and to seek for veterinarian advice if there is sign of overdosage.

Withdrawal Period (s):

Meat: 15 days.

Milk: 3 days.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure. Keep out of reach of children (Jauhkan dari kanak-kanak). Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed in accordance with local requirements.

Shelf life packaged for sale: 2 years

Shelf-life after opening: 24 hours

Packaging Available: 100 mL

Date of revision: 26-04-2021

Manufacturer:

Hanvet Pharmaceutical Company Limited (HANVET CO., LTD.)
Pho Noi A Industrial Zone, Ban Yen Nhan Town, My Hao District,
Hung Yen Province, Vietnam.

Marketing authorization holder:

NAM PHARM SDN. BHD.

5, Lebuhr Perusahan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833