

CLAVAMOX®
(amoxicillin trihydrate/
clavulanate potassium)
Drops

P1506643

23840

Name and Strength of Active Ingredients: When reconstituted each mL contains 50 mg of amoxicillin as the trihydrate and 12.5 mg of clavulanic acid as the potassium salt.

Product Description:

For dry powder: Off-white powder with bubble gum flavor.
Reconstituted Suspension: White or off-white suspension with bubble-gum flavor.

Pharmacodynamics: Amoxicillin is bactericidal in action and acts through the inhibition of biosynthesis of cell wall mucopeptide of susceptible organisms. The action of clavulanic acid extends the antimicrobial spectrum of amoxicillin to include organisms resistant to amoxicillin and other β -lactam antibiotics. Amoxicillin/clavulanate has been shown to have a wide range of activity which includes β -lactamase-producing strains of both gram-positive and gram-negative aerobes, facultative anaerobes, and obligate anaerobes. Many strains of the following organisms, including β -lactamase-producing strains, isolated from veterinary sources, were found to be susceptible to amoxicillin/clavulanate in vitro but the clinical significance of this activity has not been demonstrated for some of these organisms in animals.

Aerobic bacteria, including *Staphylococcus aureus**, β -lactamase-producing *Staphylococcus aureus** (penicillin resistant), *Staphylococcus species**, *Staphylococcus epidermidis*, *Staphylococcus intermedius*, *Streptococcus faecalis*, *Staphylococcus species**, *Corynebacterium pyogenes*, *Corynebacterium species*, *Erysipelothrix rhusiopathiae*, *Bordetella bronchiseptica*, *Escherichia coli**, *Proteus mirabilis*, *Proteus species*, *Enterobacter species*, *Klebsiella pneumoniae*, *Salmonella dublin*, *Salmonella typhimurium*, *Pasteurella multocida**, *Pasteurella haemolytica*, *Pasteurella species**.

* The susceptibility of these organisms has also been demonstrated in in vivo studies.

Studies have demonstrated that both aerobic and anaerobic flora are isolated from gingival cultures of dogs with clinical evidence of periodontal disease. Both gram-positive and gram-negative aerobic and anaerobic subgingival isolates indicate sensitivity to amoxicillin/clavulanic acid during antimicrobial susceptibility testing.

Pharmacokinetics: Clavamox is stable in the presence of gastric acid and is not significantly influenced by gastric or intestinal contents. The 2 components are rapidly absorbed resulting in amoxicillin and clavulanic acid concentrations in serum, urine and tissues similar to those produced when each is administered alone. Amoxicillin and clavulanic acid diffuse readily into most body tissues and fluids with the exception of brain and spinal fluid, which amoxicillin penetrates adequately when meninges are inflamed. Most of the amoxicillin is excreted unchanged in the urine. Clavulanic acid's penetration into spinal fluid is unknown at this time. Approximately 15% of the administered dose of clavulanic acid is excreted in the urine within the first 6 hours. Clavamox combines the distinctive properties of a broad-spectrum antibiotic and a β -lactamase inhibitors to effectively extend the antibacterial spectrum of amoxicillin to include β -lactamase-producing as well as non- β -lactamase-producing aerobic and anaerobic organisms.

Indications: Clavamox Drops are indicated in the treatment of:

Dogs: Skin and soft tissue infections such as wounds, abscesses, cellulitis, superficial/juvenile and deep pyoderma due to susceptible strains of the following organisms: β -lactamase-producing *Staphylococcus aureus*, non- β -lactamase-producing *Staphylococcus aureus*, *Staphylococcus spp.*, *Streptococcus spp.*, and *E. coli*.

Periodontal infections due to susceptible strains of both aerobic and anaerobic bacteria. Clavamox has been shown to be clinically effective for treating cases of canine periodontal disease.

Cats: Skin and soft tissue infections such as wounds, abscesses, and cellulitis/dermatitis due to susceptible strains of the following organisms: β -lactamase-producing *Staphylococcus aureus*, non- β -lactamase-producing *Staphylococcus aureus*, *Staphylococcus spp.*, *Streptococcus spp.*, *E. coli*, *Pasteurella multocida*, and *Pasteurella spp.*

Urinary tract infections (cystitis) due to susceptible strains of *E. coli*.

F000007875P11

Recommended Dose:

Add 14 mL of water to the bottle and shake vigorously.

Dogs: The recommended dosage is 6.25 mg/lb (1mL/10lb) of bodyweight twice a day. Skin and soft tissue infections such as abscesses, cellulitis, wounds, superficial/juvenile pyoderma, and periodontal infections should be treated for 5-7 days or for 48 hours after all symptoms have subsided. If no response is seen after 5 days of treatment, therapy should be discontinued and the case reevaluated. Deep pyoderma may require treatment for 21 days; the maximum duration of treatment should not exceed 30 days.

Cats: The recommended dosage is 62.5 mg (1mL) twice a day. Skin and soft tissue infections such as abscesses and cellulitis/dermatitis should be treated for 5-7 days or 48 hours after all symptoms have subsided, not to exceed 30 days. If no response is seen after 3 days of treatment, therapy should be discontinued and the case reevaluated.

Urinary tract infections may require treatment for 10-14 days or longer. The maximum duration of treatment should not exceed 30 days.

Mode of Administration: Oral route.

Contraindications: The use of this drug is contraindicated in animals with a history of an allergic reaction to any of the penicillins or cephalosporins.

Warnings and Precautions: Do not use if products is discolored.

For use in dogs and cats only.

Interactions with Other Medicaments: Before giving Clavamox, tell your veterinarian if your pet is being given another antibiotic.

Pregnancy and Lactation: Safety of use in pregnant and breeding animals has not been determined.

Adverse Effects: Clavamox contains a semisynthetic penicillin (amoxicillin) and has the potential for producing allergic reactions. If an allergic reaction occur, administer epinephrine and/or steroids.

Overdose and Treatment: Stop medication and to seek for veterinarian advice if there is sign of over dosage.

Storage Conditions: Do not store dry powder above 25°C. Refrigeration (2-8°C) is required for reconstituted suspension. Any unused portion of the reconstituted suspension must be discarded after 10 days.

Dosage Form: Oral Suspension.

Packaging: 15 mL per bottle

Manufactured by:

Aurobindo Pharma Limited
Unit-XII, Sy. No. 314, Bachuppaly Village,
Quthbullapur Mandal, R.R. District.

zoetis

Product Registration Holder :

Zoetis Malaysia Sdn. Bhd. (8898-V)

Lot 3.1, Level 3, Tower 7, Avenue 3, Bangsar South,
No. 8, Jalan Kerinchi,
59200, Kuala Lumpur, Malaysia.

Date of Revision: Feb 2016