

Erythra 5.5

Product Name: Erythra 5.5 Water Soluble Powder

Composition:

Active ingredient:

Erythromycin (as Erythromycin Thiocyanate) 5.5%w/w Water Soluble Powder

Product Description: It is a pink crystalline powder.

Pharmacodynamics

Erythromycin is a macrolide antibiotic with a bacteriostatic action against a wide range of bacteria, although it also has a bactericidal action against certain pathogenic organisms. It is highly active against gram-positive bacteria and at low concentrations it inhibits many other types of micro-organisms. In addition to the gram-positive bacteria, most strains of *Neisseria* and *Haemophilus* are sensitive as are some strains of *Bordetella*, *Brucella*, *Pasteurella*, *Listeria*, *Actinomyces*, *Mycoplasma*, *Rickettsia*, certain large viruses and *Treponema pallidum*.

Erythromycin exerts its effect only against multiplying bacteria. It inhibits protein synthesis by binding to 50 S ribosomal subunits of sensitive micro-organisms. Certain resistant micro-organisms with mutational changes in components of this subunit of the ribosome fail to bind the drug.

Pharmacokinetics

Erythromycin is absorbed rapidly when administered orally and diffusion occurs into most tissues and fluids. Elimination of erythromycin occurs primarily through hepatic metabolism and the remainder is excreted in active form in the urine and bile.

Indications:

For the treatment of Chronic Respiratory Disease caused by *Mycoplasma* infection.

Recommended Dose:

Mix 500g Erythra soluble per 107 litres (23.5 gallon) of drinking water and administer continuously for 1-5 days, depending on response (approximately 25.5 mg Erythromycin per kilogram bodyweight per day).

Slowly add 500 g to not less than 5.5 litres (1.2 gallon) of clean, cold drinking water, stirring continuously until completely dissolved. This solution should then be added to more drinking water to make up a total volume of 107 litres (23.5 gallons). Ensure that the inlet to the header tank is closed.

Contraindication

Any medicated drinking water which is not consumed within 24 hours should be discarded.

Do not use in birds producing eggs for human consumption

Warning and Precautions

i. Special precautions for use in animals

The water system to which the product is added must be in good working order and the header tanks and troughs must be free of dust, algae or other particulate matter.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Due to a lack of information on the fate and effects of erythromycin in the environment veterinary surgeons prescribing this product under the cascade should satisfy themselves that there will be no adverse impact on the environment.

ii. Special precautions to be taken by the person administering the medicinal product to animals

Do not eat, drink or smoke while preparing the diluted formulation or while animals are being treated. In case of contact with eyes, rinse immediately with clean water and seek medical advice if irritation persists. Wash hands and exposed skin after use.

Target Species

Broilers up to 6 days of age.

Birds intended as replacement layers up to 6 days of age.

Broiler-Breeders of any age.

Interaction with other Medicaments

None known

Pregnancy and Lactation

Not applicable

Side Effects

None reported

Symptoms Treatment of Overdose

No information available

Withdrawal Periods

Chickens	6 days
Eggs	6 days

Storage Condition

Store below 30°C (room temperature)

Store in a cool dry place

Any medicated drinking water not consumed within 24 hours should be discarded. At the end of the treatment, discard any unused product.

Dosage Form and Packaging Available: 227g and 1 kg

Name and address of MAH

LAI NGIT SEN
No.30, 3RD Mile Bazaar,
93200 Kuching, Sarawak.

Name and address of Manufacturer

NAM PHARMA SDN BHD
5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB
31200 Chemor, Perak, Malaysia

Date of Revision of Package Insert: 07 November 2014