



LONGAMOX® 15%W/V Suspension for Injection

COMPOSITION

Amoxicillin (as trihydrate)	15.00 g
Propylparaben	0.14 g
Excipient	Q.S.P. 100 ml

PRODUCT DESCRIPTION

A oily, white to off-white suspension for injection.

TARGET SPECIES

Cattle, pigs and lambs.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

The structure of amoxicillin includes betalactam ring and thiazolidine ring common to all penicillins.

-Lactams inhibit the formation of the bacterial cell wall by interfering with the final stage of peptidoglycan synthesis. They inhibit the transpeptidase activity of catalyzing the polymerization units of glycopeptides, the cell wall component. They have a bactericidal action only on growing cells.

The activity is exerted mainly vis-à-vis the Gram-positive bacteria, some gram negative bacteria but, in particular, Mannheimia haemolytica and Pasteurella multocida are also susceptible to the bactericidal effect of amoxicillin. Amoxicillin may be destroyed by beta-lactamases produced by some strains.

Pharmacokinetic particulars

After intramuscular administration at a dose of 15 mg / kg, amoxicillin is well absorbed with between 60 and 100% systemic bioavailability. The peak plasma concentration between 1 and 5.5 g / ml depending on the species, were observed 1-5 hours after administration.

After repeated administration (2 injections 48 hours apart), the pharmacokinetic parameters were stable and no accumulation phenomenon is observed. Plasma concentrations are maintained beyond the MIC90 for more than 32 hours after the first injection and up to 36 hours after the second injection.

Amoxicillin is active primarily eliminated by the kidney.

INDICATIONS

Diseases caused by bacteria susceptible to amoxicillin.

In cattle, pigs and lambs: treatment of respiratory infections due to Gram positive bacteria or *Pasteurella*.

CONTRAINDICATIONS

Allergy to penicillins.

Do not use in animals with known hypersensitivity to betalactams.

Do not administer to rabbits, guinea pigs, hamsters or gerbils.

Do not use in animals with severe renal impairment accompanied by anuria or oliguria.

RECOMMENDED DOSAGE

Cattle, pigs and lambs

15mg/kg (1ml/10kg bw) twice at 48 hours interval, by intramuscular route.

WITHDRAWAL PERIODS

Meat and offal:

Cattle: 35 days

Lambs: 21 days

Pigs: 27 days

Milk: 2.5 days

ADVERSE EFFECTS

Penicillins and cephalosporins may cause hypersensitivity phenomena (allergy) following administration. Allergic reactions to these substances may occasionally be severe (anaphylaxis).

INTERACTIONS WITH OTHER MEDICAMENTS

The combination of Longamox with tetracyclines, chloramphenicol and

sulphonamides alone (bacteriostatic action) is not recommended.

PREGNANCY AND LACTATION

None.

SYMPTOMS AND TREATMENT OF OVERDOSE

The trials carried out using a dose equivalent to 5 times the therapeutic dose did not induce general toxic effects.

SPECIAL PRECAUTIONS FOR STORAGE

Store the vial at a temperature below 25°C.

WARNINGS AND PRECAUTIONS

Inject a maximum volume of 20ml per injection site.

FURTHER INFORMATION

Keep out of the reach of children.

Shelf life after first opening: 1 month.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

DOSAGE FORMS AND PACKAGING

Box of one 100 ml vial

REG NUMBER : MALXXXX

FOR ANIMAL TREATMENT ONLY

CONTROLLED MEDICINE - UBAT TERKAWAL

NAME AND ADDRESS OF MANUFACTURER

VETOQUINOL S.A.
Magny-Vernois, 70200 Lure, France

NAME AND ADDRESS OF REGISTRATION HOLDER

AGE D'OR SDN BHD

15, Jalan 31/54, Section 31, Kota Kemuning 40460, Shah Alam,
Selangor D.E., Malaysia

Date of revision: December 2014

0614 H

Vétoquinol	Nom du produit : Longamox MY	
Créé par : V.MASSARD	Référence article : 0614 H	
le :	Dimensions : 160x120 mm	
Visa :	1 couleur	rev H
 281 U		