

For Animal Use Only

ETHO-480

ORAL SUSPENSION

CONTENT:

Trimethoprim80 mg/ml
Sulphadiazine.....400 mg/ml
Methyl paraben (preservative).....1 mg/ml
Propyl paraben (preservative).....0.2 mg/ml

PRODUCT DESCRIPTION: A white to light brownish viscous suspension.

PHARMACODYNAMICS:

In susceptible bacteria, the sulfonamide component blocks the synthesis of dihydrofolic acid, and the particular diaminopyrimidine used in combination inhibits the next enzyme in the sequence (dihydrofolate reductase) to prevent the formation of tetrahydrofolic acid (folinic acid). Folinic acid is required for the synthesis of DNA. This sequential blockade produces a bactericidal rather than bacteriostatic effect.

PHARMACOKINETICS:

Trimethoprim/sulfa is well absorbed after oral administration, with peak levels occurring about 1-4 hours after dosing. Trimethoprim/sulfa is well distributed in the body. When meninges are inflamed, the drugs enter the CSF in levels of about 50% those found in the serum. Both drugs cross the placenta and are distributed into milk. Trimethoprim/sulfa is both renally excreted unchanged via glomerular filtration and tubular secretion and metabolized by the liver.

INDICATION:

Bacterial infections due to susceptible organisms in chicken, turkeys and pigs.

DOSAGE & ADMINISTRATION:

Slowly add measured volume to water in medication tank, stirring thoroughly.

Infections	Inclusion Rate
<i>Escherichia coli</i> , <i>Pasteurella</i> , <i>Haemophilus</i> and <i>staphylococcal</i> , infections in chickens	1ml/4.5L water
<i>Salmonella</i> infections in chickens and diarrhoea in pigs	3ml/4.5L water
Bacterial infections in turkey	15ml/500kg flock live weight in water consumed daily

Recommended dilution rates apply when water consumption is normal; if subnormal, increase concentration to administer 15 to 30mg/kg total liveweight daily (active ingredients). Continue treatment for 3 to 5 days according to severity of disease.

CONTRAINDICATIONS:

Use in unweaned piglets less than 15 to 20 days old. Known sulfonamide sensitivity, Parenchymal hepatic damage, blood dyscrasis.

WARNINGS / PRECAUTIONS:

Should not be used in laying birds producing eggs for human consumption.

DRUGS INTERACTIONS:

Sulfadiazine solutions are incompatible with calcium or other polyionic containing fluid. Antacid tends to inhibit the GI absorption of sulfadiazine.

PREGNANCY AND LACTATION: None known.

ADVERSE EFFECTS:

Side effects due to the potentiated sulfonamides are quite rare, although adverse reactions to the sulfonamide components still occur. Up to 10 times the recommended dose of trimethoprim has been given with no adverse effects. Prolonged administration of trimethoprim at reasonably high levels leads to maturation defects in hematopoiesis due to impaired folinic acid synthesis. This effect is readily reversible by supplementation with folinic acid.

OVERDOSE AND TREATMENT: Stop medication and to seek for veterinarian advice.

STORAGE CONDITIONS: Store below 30°C.

WITHDRAWAL PERIOD(S):

Poultry and Swine

Meat: 14 days

Eggs: Do not use in birds producing egg for human consumption.

DOSAGE FORMS AND PACKAGING AVAILABLE: 1.0 liter Oral Suspension packed into HDPE bottle.

DATE OF REVISION OF PACKAGE INSERT: 21 January 2015.

Manufactured by:

PAHANG PHARMACY SDN. BHD. (10826-K)

LOT 5979, JALAN TERATAI,

5½ MILE OFF JALAN MERU, 41050 KLANG,

SELANGOR D.E., MALAYSIA.