

[Draft of Package Insert]

For Animal Use Only

TRI-SQ

WATER SOLUBLE POWDER

CONTENT: Each gram contains:

Sulfaquinoxaline500 mg

Trimethoprim.....165 mg

PRODUCT DESCRIPTION: Fine Homogenous light yellowish powder.

PHARMACODYNAMICS:

Trimethoprim is a dihydrofolate reductase inhibitor, effectively against a wide range of bacteria, including *E. Coli* and some *Klebsiella*, *Proteus* and *Staphylococcus spp*. It is used in combination with sulfonamides to prevent the conversion of bacterial dihydrofolate acid to tetrahydrofolic acid and consequently affect the DNA synthesis. Sulfaquinoxaline is a potent inhibitor of vitamin K epoxide reductase. Sulfaquinoxaline has been utilized alone or in combination with Trimethoprim for control of coccidia, colisepticemia and pasteurellosis in poultry, and some susceptible bacterial diseases.

PHARMACOKINETICS:

Sulfaquinoxaline and Trimethoprim are both rapidly absorbed from the gastrointestinal tract when administered orally.

Sulfonamides are widely distributed throughout the body into soft tissues. The ability binding to plasma proteins are varies from sulfonamide to sulfonamide and from species to species. Sulfaquinoxaline is enteric sulfonamide primarily eliminated via the feces. Trimethoprim is widely distributed to various tissues and fluids including kidneys, liver, lungs and bronchial secretions, saliva, aqueous humour, prostatic tissue and fluid, and vaginal secretions. Trimethoprim is excreted primarily by the kidneys.

INDICATION:

TRI-SQ is recommended for the treatment of colisepticaemia in chicken and turkey and for the control of coccidiosis in chickens.

The product is indicated for use in drinking water only for the treatment of coccidiosis in chickens and for the treatment of colisepticaemia in chicken and turkey.

DOSAGE & ADMINISTRATION:

Dosage:

The recommended dose is the equivalent of 30 mg combined trimethoprim and sulphaquinoxaline per kg bodyweight daily. This dose is provided by the following inclusion rates:

In drinking water: 28 g per 100 litres (equivalent to 4 sachet of TRI-SQ)

Treatment should be repeated daily for 5 days or until 2 days after symptoms have subsided, but not exceeding 14 days in total.

Administration:In drinking water:

Make an initial TRI-SQ at the rate of 28 g granules in 2 litres of cold water.

This should be stirred until well suspended. Then should be added to the poultry drinker or water header tank at approximately the rate of 2 litres to 98 litres of drinking water.

The mixture should be gently stirred until the powder has dissolved.

TRI-SQ will retain its potency for 24 hours as a solution in drinking water. It should always be prepared shortly before use. Do not use in water-proportioner systems.

Remove all other sources of water during the entire period of treatment.

CONTRAINDICATIONS:

Do not use in case of renal or hepatic impairment, sulphonamide hypersensitivity or blood dyscrasias.

Do not use in layers and breeders, birds producing eggs for human consumption.

Do not use in turkeys less than 21 days of age.

Do not use together with coccidiostats and other medications that contain sulphonamides.

WARNINGS / PRECAUTIONS:

Do not use in water proportioner system.

Do not use as a long term treatment that exceed 14 days.

Animal must not slaughtered for human consumption during treatment.

Handling this product with care to avoid direct skin contact and inhalation of any dust.

Wash hands and exposed skin with soap and water after use.

DRUGS INTERACTIONS: None known.

PREGNANCY AND LACTATION: None known.

ADVERSE EFFECTS: None known.

OVERDOSE AND TREATMENT:

A decreased in appetite and water consumption, lowered egg production, egg weight, and hatchability were reported from the dosage higher than recommended concentration.

STORAGE CONDITIONS: Store below 30°C.

WITHDRAWAL PERIOD(S):

Chickens: 10 days from the last treatment

Turkeys: 14 days from the last treatment

DOSAGE FORMS AND PACKAGING AVAILABLE: TRI-SQ is supplied in 112g aluminum sachet.

DATE OF REVISION OF PACKAGE INSERT: 18 December 2014.

Manufactured by:

PAHANG PHARMACY SDN. BHD. (10826-K)

LOT 5979, JALAN TERATAI,
5½ MILE OFF JALAN MERU, 41050 KLANG,
SELANGOR D.E., MALAYSIA.

Product Registration Holder:

GENPHARMA SDN. BHD. (268066-W)

LOT 5979, JALAN TERATAI,
5½ MILE OFF JALAN MERU, 41050 KLANG,
SELANGOR D.E., MALAYSIA.