

TM-400* gram/kg powder

OXYTETRACYCLINE DIHYDRATE 40% POWDER

FORMULA:

Each kilogram of TM-400 contains Oxytetracycline Dihydrate equivalent to the amount of 400 grams Oxytetracycline Hydrochloride, rice hulls (carrier) q.s.ad.1kg

Active Drug Ingredient: Oxytetracycline Dihydrate equivalent to Oxytetracycline Hydrochloride.

PRODUCT DESCRIPTION:

Buff Colored, free flowing powder, free from agglomerates

PHARMACODYNAMICS:

Oxytetracycline is a broad-spectrum antibiotic. Its mode of action is bacteriostatic by inhibition of prokaryotic protein synthesis at the ribosomal level. It is active against Gram-positive and Gram-negative aerobic and anaerobic micro-organisms, *Mycoplasma* spp., *Chlamydia* spp., *Leptospira* spp. and Rickettsias. It has little activity against *Salmonella* spp., *Escherichia coli*, Enterobacter and *Klebsiella* spp. and has essentially no activity against *Pseudomonas* spp., *Proteus* spp. and yeasts. Sub-Minimum Inhibitory Concentration (MIC) effects have been identified for tetracyclines. Inhibitions of adhesion of pathogens to host cell surfaces are known to occur at levels of antibiotic lower than the MIC. Host cell adhesion is an essential pre-requisite for the expression of pathogenicity for several porcine respiratory pathogens e.g. *Actinobacillus pleuropneumoniae*, *Bordetella bronchiseptica* and mycoplasmas. Selected resistance to tetracyclines is not uncommon among bacteria and it is important that sensitivity is determined prior to initiation of therapy. Most organisms which are resistant to one member of the tetracycline group will have cross-resistance to other members of the group.

PHARMACOKINETICS:

Due to the amphoteric nature of Oxytetracycline, oral bioavailability is low (<10%). Oxytetracycline has excellent distribution properties with a Volume of Distribution (Vd) exceeding 1.2 L/kg. The compound tends to concentrate in the lung, liver and kidney. Oxytetracycline undergoes enterohepatic circulation. It does not appear to be metabolised to any significant degree. The compound is mainly excreted as the parent molecule primarily in the urine but also in the faeces.

ENVIRONMENTAL PROPERTIES:

The same as many therapeutic antibiotics used in animal feed, tetracycline compounds can affect microorganisms in the environment if not properly disposed. Care should be taken to assure that leftover feeds with TM400 and empty packages of the product are properly disposed of in accordance with local authorities regulations.

INDICATIONS AND RECOMMENDED DOSAGES:

Chickens:

Amount: 100 -200 grams Oxytetracycline per ton of feed.

- Control of infectious synovitis caused by *Mycoplasma synoviae*; control of fowl cholera caused by *Pasteurella multocida* susceptible to oxytetracycline. Feed continuously for 7 to 14 days.

Amount: 400 grams Oxytetracycline per ton of feed.

- For control of chronic respiratory disease (CRD) and air sac infection caused by *M. gallisepticum* and *Escherichia coli* susceptible to oxytetracycline. Feed continuously for 7 to 14 days.

Amount: 500 grams Oxytetracycline per ton of feed.

- For reduction of mortality due to air sacculitis (air-sac infection) caused by *E. coli* susceptible to oxytetracycline. Feed continuously for 5 days.

Swine:

Amount: 10 milligrams Oxytetracycline per pound of body weight.

- For treatment of bacterial enteritis caused by *E. coli* and *Salmonella choleraesuis* susceptible to oxytetracycline and treatment of bacterial pneumonia caused by *P. multocida* susceptible to oxytetracycline. Feed continuously for 7 to 14 days.

Swine (Breeding):

Amount: 10 milligrams Oxytetracycline per pound of body weight.

- For control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospiras) caused by *Leptospira Pomona* susceptible to oxytetracycline. Feed continuously for not more than 14 days.

MODE OF ADMINISTRATION:

Route of Administration: ORAL

Mixing Directions: Determine the appropriate amount of TM-400 required from the recommended dosage. Mix thoroughly in feed.

CONTRAINDICATIONS:

Oxytetracycline in low calcium feed (0.18 - 0.55% dietary calcium) should not be fed for more than 5 days and should not be fed to laying hens.

WARNINGS AND PRECAUTIONS:**Special precautions for use in animals**

- Long term use of this product may lead to development of bacterial resistance and is not recommended. However, the recommended course of treatment should be completed.
- For use in dry animal feed only. Not for use in liquid feed supplements.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

- If allergic to oxytetracycline, do not handle the product.
- When incorporating into feed, care should be taken not to inhale any dust when handling the product and skin contact should be avoided.
- It is recommended that a face mask, be worn during the dispensing of the product.
- Hands and exposed skin should be washed thoroughly at the end of the operation.

INTERACTIONS WITH OTHER MEDICAMENTS:

Tetracyclines may be partially bound and inactivated when included in feeds with high content of calcium salts.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Do not administer to chickens producing eggs for human consumption

ADVERSE EFFECTS / UNDESIRABLE EFFECTS:

No known side effects are expected when directions are followed.

OVERDOSE AND TREATMENT:

If overdose is confirmed, feed should be removed.

WITHDRAWAL PERIOD(S):**Chickens:**

24-hour withdrawal period at 500 g/ton use level.

0 (zero) days at lower use levels.

3 days before slaughter in low calcium feeds.

Swine:

0 (zero) days before slaughter at 10 milligrams per pound.

.

STORAGE CONDITIONS:

Stored in dry place below 30°C.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Medicated powder premix

25 kg bag

Manufactured by Range Pharma Sdn Bhd

No. 1, Jln TSB 11, Taman Industri Sg. Buloh, 47000 Sungai Buloh, Selangor, Malaysia.

Registration holder: Phibro Corporation (M) Sdn Bhd, 311, Level 3, Menara Axis, 2, Jalan 51A/223, Seksyen 51A, 46100 Petaling Jaya, Selangor, Malaysia

Revision Date of Package Insert: Aug 15, 2025