

TM-200* gram/kg powder

BROAD SPECTRUM ANTIBIOTIC CONTAINING OXYTETRACYCLINE DIHYDRATE 20%

FORMULA:

Each kilogram of TM-200 contains Oxytetracycline Dihydrate equivalent to the amount of 200 grams Oxytetracycline Hydrochloride, rice hulls (carrier) q.s.ad.1kg

Active Drug Ingredient: Oxytetracycline Dihydrate equivalent to Oxytetracycline Hydrochloride.

PRODUCT DESCRIPTION:

Buff Colored, free flowing powder, free from agglomerates

PHARMACODYNAMICS:

Oxytetracycline (Dihydrate) is a naturally occurring tetracycline, produced by *Streptomyces rimosus*. Like other tetracycline it has a broad spectrum of activity. It was introduced in 1950. Effective against a wide range of both gram positive and gram negative organisms. The main mechanism of action is on protein synthesis.

PHARMACOKINETICS:

Oxytetracycline is absorbed from the gastro-intestinal tract, and under high dosage, it is widely distributed in tissues and fluids and is excreted in bile, stool, and urine.

INDICATIONS AND RECOMMENDED DOSAGES:

Chickens:

Amount: 100 -200 grams Oxytetracycline per ton of feed.

- Control of infectious synovitis caused by *Mycoplasma synoviae*; control of fowl cholera caused by *Pasteurella multocida* susceptible to oxytetracycline. Feed continuously for 7 to 14 days.

Amount: 400 grams Oxytetracycline per ton of feed.

- For control of chronic respiratory disease (CRD) and air sac infection caused by *M. gallisepticum* and *Escherichia coli* susceptible to oxytetracycline. Feed continuously for 7 to 14 days.

Amount: 500 grams Oxytetracycline per ton of feed.

- For reduction of mortality due to air sacculitis (air-sac infection) caused by *E. coli* susceptible to oxytetracycline. Feed continuously for 5 days.

Swine:

Amount: 10 milligrams Oxytetracycline per pound of body weight.

- For treatment of bacterial enteritis caused by *E. coli* and *Salmonella choleraesuis* susceptible to oxytetracycline and treatment of bacterial pneumonia caused by *P. multocida* susceptible to oxytetracycline. Feed continuously for 7 to 14 days.

Swine (Breeding):

Amount: 10 milligrams Oxytetracycline per pound of body weight.

- For control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospiras) caused by *Leptospira Pomona* susceptible to oxytetracycline. Feed continuously for not more than 14 days.

MODE OF ADMINISTRATION:

Orally via feed. Mixing Directions: Determine the appropriate amount of TM-200 required from the recommended dosage. Mix thoroughly in feed.

CONTRAINDICATIONS:

Oxytetracycline in low calcium feed (0.18 - 0.55% dietary calcium) should not be feed for more than 5 days

WARNINGS AND PRECAUTIONS:

Under veterinarian recommendation.

INTERACTIONS WITH OTHER MEDICAMENTS:

Tetracyclines may be partially bound and inactivated when included in feeds with high content of calcium salts.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Do not administer to chickens producing eggs for human consumption.

ADVERSE EFFECTS / UNDESIRABLE EFFECTS:

The severe or irreversible adverse effects of Oxytetracycline (Dihydrate), which give rise to further complications include renal failure.

OVERDOSE AND TREATMENT:

If overdose is confirmed, feed should be removed.

WITHDRAWAL PERIOD(S):**Chickens:**

24-hour withdrawal period at 500 g/ton use level.

0 (zero) days at lower use levels.

3 days before slaughter in low calcium feeds.

Swine:

0 (zero) days before slaughter at 10 milligrams per pound.

STORAGE CONDITIONS:

Store in dry place below 30°C.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Medicated powder premix

25 kg bag

Manufactured by Range Pharma Sdn Bhd

No. 1, Jln TSB 11, Taman Industri Sg. Buloh, 47000 Sungai Buloh, Selangor, Malaysia.

Registration holder: Phibro Corporation (M) Sdn Bhd

311, Level 3, Menara Axis,

2, Jalan 51A/223, Seksyen 51A,

46100 Petaling Jaya, Selangor, Malaysia

Revision Date of Package Insert: Aug 15, 2025