

Amoxy 500 mg/g
Soluble Powder
(FOR VETERINARY USE ONLY)

COMPOSITION

Amoxicillin trihydrate 500 mg/g

PRODUCT DESCRIPTION

This product is fine white color powder.

TARGET SPECIES

Chickens, Turkeys, Ducks and Pigs

PHARMACODYNAMICS

Amoxicillin is a bactericidal antibiotic belonging to the group of semisynthetic penicillins with a broad spectrum of activity against Gram positive and Gram negative bacteria. It owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

PHARMACOKINETICS

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids. Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared to be a more important route of elimination in birds than in mammals.

INDICATIONS

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

Pigs: For the treatment of pasteurellosis.

CONTRA-INDICATIONS

This product should not be administered to rabbits, guinea pigs, hamsters, gerbils or any other small herbivore.

Do not use in animals with known hypersensitivity to penicillins or other β -lactam antibiotics or to any of the excipients.

Do not administer to animals with renal disease including anuria or oliguria.

ADMINISTRATION & DOSAGE

In drinking water use.

Chickens: The recommended dosage is 15 mg amoxicillin trihydrate per kg bodyweight per day (corresponding to 30 mg product/kg-bodyweight/day). The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks: Recommended dosage is 20 mg amoxicillin trihydrate/kg bodyweight per day (corresponding to 40 mg product/kg bodyweight/day) for 3 consecutive days.

Turkeys: Recommended dosage is 15-20 mg amoxicillin trihydrate/kg bodyweight per day (corresponding to 30-40 mg product/kg-bodyweight/day) for 3 days or in severe cases for 5 days.

Pigs: Administer in the drinking water to give 20 mg amoxicillin trihydrate/kg bodyweight (corresponding to 40 mg product/kg-bodyweight) daily for up to 5 days.

The following formula may be used to calculate the required concentration of product (in milligrams of product per liter of drinking water):

x mg product per kg bodyweight per day	$\times$	mean bodyweight (kg) of animals to be treated	=	x mg product per litre drinking water
mean daily water consumption (l) per animal				

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing. The uptake of medicated water depends on the clinical condition of the animal. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake.

After the end of the medication period the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

WITHDRAWAL PERIODS

Meat and offal:

- Chickens: 1 day
- Ducks: 9 days
- Turkeys: 5 days
- Pigs: 2 days

Not authorised for use in laying birds producing eggs for human consumption and within 3 weeks of onset of laying.

WARNING AND PRECAUTIONS

Special warnings for each target species:

None

Special precautions for use:

i) Special precautions for use in animals:

Not effective against beta-lactamase producing organisms. Pigs: The uptake of medication by animals may be altered as a consequence of illness. In case of insufficient water uptake, animals should be treated parenterally instead. Official, national and regional antimicrobial policies should be taken into account when the product is used. Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria. Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the amoxicillin and may decrease the effectiveness of the treatment.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious. People with known hypersensitivity to the active substance or if you have been advised not to work with such preparations should avoid contact with the veterinary medicinal product. Handle this product with great care to avoid exposure, taking all recommended precautions. If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms

and require urgent medical attention. Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Wear gloves during preparation and administration of medicated water or liquid feed. Wash any exposed skin after handling the product or medicated water or feed. Wash hands after use.

- For Veterinary Use Only
- Keep out of reach of children.
- Jauhi ubat dari kanak-kanak.

INTERACTION WITH OTHER MEDICAMENTS

The product should not be administered with antibiotics that have a bacteriostatic mode of action, such as tetracyclines, macrolides, sulphonamides.

PREGNANCY AND LACTATION

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin. Use only according to the benefit/risk assessment of the responsible veterinarian. Do not use in birds producing eggs for human consumption or within 3 weeks of onset of laying.

SIDE EFFECT

Penicillins and cephalosporins may cause hypersensitivity reactions which may occasionally be serious.

OVERDOSE AND TREATMENT

No problems with overdose have been reported. Treatment should be symptomatic and no specific antidote is available.

SHELF LIFE: 36 months

SHELF LIFE after first opening: 24 hours

SHELF LIFE after dilution: 12 hours

STORAGE

Store in a cool dry place ($\leq 25^{\circ}\text{C}$). Strictly avoid light & heat exposure

PACKAGING

The powder filled 100 g in aluminum sachets, by automatic filling machine and sealing, and packed in carton box.

DATE OF REVISION: 10 Jun 2019



Manufactured & Distributed By:

SHENNONG ANIMAL HEALTH (M) SDN BHD
神農動物保健(馬)有限公司
No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,
Taman Perindustrian USJ 1,
47610 Subang Jaya, Selangor.