

Veterinary Use Only

PENTASOL

Amoxicillin 50% w/w Water Soluble Powder

1. Brand or Product Name/ Name and Strength of Active Substance(s)

Pentosol Amoxicillin 50% w/w Water Soluble Powder

Each kg contains:

Amoxicillin (as amoxicillin trihydrate) ----- 500g

2. Product Description

White or pale yellow powder

3. Pharmacodynamics / Pharmacokinetics / Environmental Properties**Pharmacodynamics :**

Amoxicillin is a semi-synthetic, acid-stable broad spectrum penicillin. It inhibits peptidoglycan synthesis of the bacterial cell wall and has a fast bacteriocidal action. Amoxicillin is active against a number of Gram-positive and Gram-negative bacteria.

Pharmacokinetics :

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids. Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

4. Indication

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

Not effective against beta-lactamase producing organisms.

Use of the product should be based on susceptibility testing and it should take into account official and local antimicrobial policies.

Pigs : For the treatment of salmonellosis and pasteurellosis

5. Recommended Dosage/ Mode of Administration

In drinking water use/ in feed use.

Chickens, turkeys and ducks :

The product is administered in the drinking water.

Prepare the solution with fresh tap water immediately before use.

Once opened, use the contents of the package immediately.

Any unused medicated water should be discarded after 12 hours.

Chicken : 13 mg amoxicillin (as trihydrate) per kg body weight.

The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks : 17 mg amoxicillin (as trihydrate) per kg body weight for 3 consecutive days.

Turkeys : 13-17 mg amoxicillin (as trihydrate) per kg body weight for 3 days or in severe cases for 5 days

Pigs : For the medication of Pigs, the product may be administered in the drinking water or administered by addition to liquid feeds produced with commercial feed
It may not be used in dry feeds.

1. Administration via the drinking water Administer in the drinking water to give 17mg/kg bodyweight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days.

2. Administration in liquid feed Administer in the liquid feed, to give 17 mg amoxicillin (as trihydrate)/kg bodyweight daily for up to 5 days. Medicated feed should be freshly prepared on at least 3 occasions per day over the treatment period.

The daily dose should be calculated based on the number of animals and average weight and then divided by the number of feed lots prepared in the day.

6. Contraindications

- Do not use in animals with known hypersensitivity to penicillins or other substances of the β -lactam group.
- Do not use in animals with serious kidney malfunction including anuria and oliguria.
- Do not use in presence of β - lactamase producing bacteria.
- Should not be administered to rabbits, hamsters, gerbils and guinea pigs, or to birds producing eggs intended for human consumption.

7. Warnings and Precautions

- Use under veterinarian recommendation.
- Persons who are allergic to penicillins and cephalosporins should avoid direct contact with the product.
- Keep out of reach of children/Jauhi ubat daripada kanak-kanak.

8. Interactions with Other Medicaments

None known.

9. Statement on usage during pregnancy and lactation

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

10. Adverse Effects/Undesirable Effects

- Allergic reactions seldom may occur.
- Gastrointestinal side effects (diarrhoea) possibly may occur.

11. Overdose and Treatment

No problems with over dosage have been reported. Treatment should be symptomatic and no specific antidote is available.

12. Withdrawal Period(s)

Meat & offal

Chickens : 1 day

Ducks : 9 days

Turkeys : 5 days

Pigs : 2 days

Not for use in laying birds producing eggs for human consumption.

13. Shelf life (after reconstitution or dilution)

3 hours for liquid feed and 12 hours in drinking water after dilution

14. Storage Condition

Store in a dark and cool place with dry condition below 30°C.

15. Dosage Forms and packaging available

1kg

16. Name and Address of manufacturer/marketing authorization holder

Nutri Pharmax Sdn Bhd

6, Lorong Serunai 1, Valdor Sungai Bakap, 14200 Sungai Jawi, Pulau Pinang, Malaysia.

17. Date of Revision of Package Insert

08 April 2020



Manufactured by

KBNP, INC.

235-9 Chusaro, Sinam, Yesan,
Chungnam, Korea



Distributed by

Nutri Pharmax Sdn.Bhd.

6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang, Malaysia
Tel: +604-582 9377