

CTC 500 Powder

Brand or Product Name:

CTC 500 Powder

Name and Strength of Active Substance(s)

Each g of CTC 500 powder contains Chlortetracycline Hydrochloride 500mg (strength 50% w/w).

Each Kg of CTC 500 powder contains Chlortetracycline Hydrochloride 500g (strength 50% w/w).

Product Description

CTC 500 powder is a yellow powder packed in 500g aluminum sachet or 25kg paper bag

Pharmacodynamics/Pharmacokinetics/Environmental Properties

Chlortetracycline is used as an antimicrobial in veterinary therapeutics. As with other tetracyclines, it is a broad spectrum antibiotic with a wide range of bacteriostatic activity against both gram negative and gram positive bacteria. Chlortetracycline appears to act by inhibiting the binding of aminoacyl-tRNA to the mRNA ribosome complex, thereby preventing the prokaryotic cell's ability to synthesise proteins.

Following oral administration of a single dose, chlortetracycline is absorbed from the stomach and the first part of the small intestine to give a peak plasma concentration in 2 - 4 hours. A gradual fall in level follows: at 24 hours, the substance is barely detectable. Some of the drug is concentrated by the liver, excreted in bile and resorbed from the intestine; a small amount persists in the blood as a result of enterohepatic resorption.

Indication:

Pigs: As an aid in the control of diseases caused by chlortetracycline-sensitive organisms in swine

Chickens: The product is effective in the treatment and control of E. coli septicaemia secondary to Infectious Bursal Disease (Gumboro Disease), Chronic Respiratory disease (CRD) caused by Mycoplasma gallisepticum and E. coli and Salmonella enteritidis, Salmonella typhimurium, Moraxella anatipestifer and Pasteurella multocida infection.

Turkeys: For the treatment and control of diseases responsive to chlortetracycline, including secondary E.coli infections involved in outbreaks of Turkey Rhinotracheitis (TRT).

Targeted Species: Pigs, chickens and turkeys**Recommended Dosage:**

Pigs: For oral administration following incorporation into the complete feed to achieve a dose rate of 20mg chlortetracycline hydrochloride per kg live weight daily.

Chickens: Recommended dosage is 20 - 30mg chlortetracycline hydrochloride per kg bodyweight daily.

The recommended level of chlortetracycline hydrochloride in feed is generally achieved at 300 – 400ppm, that is by mixing the product into the feed at a rate of 600 – 800g per tonne of feed.

Treatment should be given for 5 - 7 days for pigs and chickens.

Turkeys: For oral administration into the complete feed to achieve a dose rate of 20mg chlortetracycline hydrochloride per kg live weight daily. This may be achieved by the incorporation of 800g – 1.2kg product per tonne complete feed. Treat for 10 consecutive days.

Feed containing the product should constitute the entire feed provided for animals and birds during treatment. Treatment should be maintained until signs of disease are no longer apparent.

Mode of Administration:

CTC 500 is to be mixed homogeneously before administration

Contraindications

Do not administer to ruminants: if administered to ruminating animals, toxicity is likely to be seen.

Warnings and Precautions

If you know you are hypersensitive (allergic) to chlortetracycline, do not handle the product. When handling this product inhalation of the dust must be avoided by wearing a disposable half-mask respirator. Avoid contact with skin and eyes. Gloves should be worn whilst handling this product. If contact with skin or eyes occurs, wash area immediately with copious amounts of fresh water. If irritation persists seek medical attention. Do not eat, drink or smoke whilst handling the product. Hands and exposed skin should be washed thoroughly after use.

Interactions with Other medications

Synergism between tetracyclines and tylosin against *Pasteurella* has been described and may occur with other macrolides and other bacteria. Combination with polymyxins may also give synergistic effects by enhancing bacterial uptake of the drugs

Pregnancy & Lactation: None known.

Adverse Effects / Undesirable Effects

From a pharmacologic viewpoint, the tetracyclines are relatively safe drugs. Adverse effects may be attributed to their severely irritant nature (vomiting after oral administration), its disturbance of intestinal flora, its ability to bind calcium and its toxic effects on liver and kidney cells

Overdose and Treatment

In case of overdosage, discontinue medication, treat symptomatically and institute supportive measures..

Withdrawal Period(s)

Animals and birds must not be slaughtered for human consumption during treatment. Pigs may only be slaughtered for human consumption 7 days after the last treatment. Chickens may only be slaughtered for human consumption 3 days after the last treatment. Turkeys may only be slaughtered for human consumption 1 day after the last treatment. This product must not be used in birds producing eggs for human consumption

Shelf Life

3 years from date of production. See disposal of containers for opened package.

Maximum Residual Limits (MRL)

Pigs: Muscle: 200µg/kg
Liver: 600µg/kg
Kidney: 1200µg/kg

Poultry: Muscle: 200µg/kg
Liver: 600µg/kg
Kidney: 1200µg/kg
Eggs: 400µg/kg

Ref. REGOV, Version 2, dated Aug 2008]

Storage Conditions

Store in a dry place below 25°C. Keep out of reach from children.

Disposal of containers

Dispose used containers in trash.

An opened package or unused medicated feed is to be discarded after 24 hours.

Dosage Forms and packaging available

Available in 500g Aluminum sachet or 25Kg paper bag packing

Name and Address of Marketing Authorization Holder:

Life (Veterinary) Sdn. Bhd.

No. 115, Taman AST, Jalan Sungai Ujong, 70200 Seremban, Negeri Sembilan, Malaysia

Name and Address of manufacturer:

Life Biopharma Sdn. Bhd.

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia

Date of Revision of Package Insert: 12-June 2015