

AMOX-500MG PREMIX

FOR ANIMAL USE ONLY

Product Name

AMOX-500MG PREMIX

Name and Strength of Active Substance(s)

Each kilogram contains 500g Amoxicillin which is equivalent to 574g Amoxicillin Trihydrate.

Product Description

AMOX-500MG Premix is a white to off white powder.

Pharmacodynamics Properties

Amoxicillin is a bactericidal antibiotic of the beta-lactam family which acts by blocking the biosynthesis of the cell wall. As a bactericidal antibiotic, amoxicillin inhibits the development of the peptidoglycan network structure of the bacterial cell wall. It is thought to act on transpeptidation which is the last step in the synthesis of the final bacterial-membrane structure.

Pharmacokinetics Properties

After oral administration in feed, amoxicillin bioavailability is approximately 30%. The fraction reaching the blood stream is metabolized to a slight extent. The metabolite identified in laboratory species is penicilloic acid that is rapidly eliminated in the urine. Amoxicillin is weakly bound to plasma proteins (about 20%) and is widely distributed in different organs.

Indication

For the control of diseases caused by *Streptococcus suis* in weaned piglets.

Recommended Dosage

30mg of AMOX-500MG PREMIX (equivalent to 15 mg of amoxicillin) per kg bodyweight daily for 14 consecutive days.

This dose may be achieved by the addition of :

- 1) 400 ppm or 800g of AMOX-500MG PREMIX (equivalent to 400 g of amoxicillin) per ton of feed in starter feed.
 - 2) 300 ppm or 600g of AMOX-500MG PREMIX (equivalent to 300 g of amoxicillin) per ton of feed in feed intended for piglets older than 6 weeks.
- To ensure thorough dispersion, the product should first be mixed with a suitable quantity of feed before incorporation in the final mix.

Mode of Administration

Oral use. To be administered in feed.

Contraindications

Do not use in animals with serious kidney (accompanied by anuria and oliguria) and/or liver failure.

Do not use in animals with known hypersensitivity to beta-lactams.

The use of the product is contraindicated when a resistance to amoxicillin is known.

Do not use in the presence of β -lactamase producing bacteria.

Do not administer to rabbits, guinea-pigs, hamsters or gerbils.

Warnings and Precautions

Special warnings

Animals with reduced feed intake and/or disturbed general condition have to be treated parenterally.

Special precautions for use

(i) Special precautions for use in animals

Animals with distinctive clinical signs of streptococcosis require individual treatment. Use of the veterinary medicinal product should be based on susceptibility testing and take into account official and local antimicrobial policies. Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to amoxicillin.

(ii) Special precautions to be taken by person handling the product :

When feed is being prepared, avoid skin contact. Whilst handling the product, wear a mask, coveralls, protective goggles and gloves at all times. Avoid inhalation of dust. Do not smoke or eat during use. Do not handle this product if you are sensitive to penicillins. Wash hands after use.

Interactions with Other Medicaments

Do not administer together with bacteriostatic anti-infectious agents (tetracyclines, sulphonamids)

Do not use simultaneously with neomycin as it prevents the absorption of oral penicillins

Pregnancy and lactation

Laboratory studies have not produced any evidence of teratogenic, embryotoxic or maternotoxic effects of amoxicillin. Nevertheless, no study was performed in the target species during pregnancy or lactation. Therefore, use only accordingly to the benefit/risk assessment by the responsible veterinarian.

Adverse Effects

Gastrointestinal signs may sometimes be observed (diarrhea). Penicillins may cause allergic reactions after the administration of the product. Allergic reactions to these substances may sometimes be severe (anaphylaxis).

Symptoms and treatment of Overdose

Penicillins have a wide margin of safety.

Withdrawal Period(s)

Meat : 4 days

Shelf-life

Shelf-life of the veterinary medicinal product as packaged for sale: 2 year.

Shelf life after first opening and incorporation into feed: 24 hours.

Allowable Maximum Residual Limit (MRL)

All food producing species : Muscle, liver, kidney, fat = 50 μ g/kg [Ref. REGOVP, Version 3-July 2014]

Storage Conditions

Store in tightly-closed original container, in a dry place below 30°C. Keep away from children. Jauhkan ubat daripada kanak-kanak.

Disposal of containers

Empty bags can be safely disposed in trash according to local government regulations.

Packaging

Net 25 kg powder packed in a layer of low density polyethylene plastic bag within a layer of polypropylene-lined paper bag.

Product Registration Holder and Manufactured by

Life Biopharma Sdn. Bhd. (546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Last Revision of Package Insert

2015-11-13