

Avadol®-Fast Tablet 500mg (Active ingredient: Paracetamol 500mg)

Product Description: Avadol®-Fast tablet is a white to off-white, round tablet with 500 debossed on one side and plain on the other side.

Pharmacodynamics: Paracetamol is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties.

Analgesic: The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (specifically cyclooxygenase (COX)-2) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation.

The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation

Antipyretic: Paracetamol act centrally on the hypothalamic heat-regulating center to produce peripheral vasodilatation resulting in increased blood flow through the skin, sweating and heat loss. Paracetamol reduces fever by inhibiting the formulation and release of prostaglandins in the CNS and by inhibiting endogenous pyrogens at the hypothalamic thermoregulator center.

Pharmacokinetics: Following oral administration paracetamol is rapidly absorbed. Paracetamol absorption takes place mainly in the small intestine and therefore the rate of absorption is depending on the rate of gastric emptying. It has been shown that drugs which delay gastric emptying also delay the absorption of paracetamol whereas metoclopramide (a drug which increases the rate of gastric emptying) accelerates absorption of the analgesic though the total amount absorbed does not increase.

The presence of food in the stomach has also been reported to reduce the rate of absorption of paracetamol. Alterations in gastric pH have no appreciable effect on paracetamol absorption. During absorption, the amount of paracetamol which is inactivated is negligible and it has been shown that paracetamol does not affect gastric mucosal permeability and does not produce mucosal bleeding. Peak plasma concentrations are reached 1 hour after absorption. The plasma half-life is 1 to 3 hours. Paracetamol penetrates the brain and is present in breast milk of human. Paracetamol is metabolized by the microsomal enzyme system of the liver. This metabolism is mainly to the glucuronide and sulphate conjugates, accounting for approximately 49% and 26% of the ingested dose respectively. About 4% is excreted as free paracetamol. Other minor pathways include the production of catechol derivatives and cysteine conjugates (via glutathione). Paracetamol excretion is rapid and occurs via the urine.

Indication For the relief of fever, mild to moderate pain including: headache, migraine, backache, musculoskeletal pain, myalgia, dysmenorrhea (period pain), pain of osteoarthritis, toothache, and the discomfort from colds, influenza and sore throats.

Recommended Dosage: Adults: 1-2 tabs for 3 to 4 times a day. Maximum: 8 tablets /day. Not recommended for children.

Mode of Administration: Oral

Contraindications: Hypersensitivity to paracetamol or any of the other ingredients/components of the product. Taking other medicines containing paracetamol. Severe and active hepatic impairment

Warnings and Precautions

This preparation contains PARACETAMOL.
Do not take any other paracetamol
containing medicines at the same time.

- Keep out of reach of children.
- Do not take if allergic to paracetamol.
- Patients should contact their health care provider if symptoms persist (if the pain lasts for more than 10 days, if there is redness or fever lasting more than 3 days).
- Paracetamol should be given with care to patients with alcohol dependence, impaired kidney or liver function.
- Large doses should be avoided in patients with hepatic impairment. Paracetamol overdose may harm the liver.
- Do not exceed recommended dose.

• Paracetamol provides symptomatic relief only, additional therapy to treat the cause of the pain or fever should be instituted when necessary.

• **Allergy alert:** Paracetamol may cause severe skin reactions, symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Effects on Ability to Drive and Use Machines

It is unlikely to impair a patient's ability to drive or use machinery.

Interactions with Other Medicaments: The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Statement on Usage During Pregnancy and Lactation

Use in pregnancy:

- Considered to be the analgesic of choice in pregnant patients.
- Although it crosses placenta, paracetamol is considered to be safe in normal therapeutic doses for short-term use as a minor analgesic/antipyretic in pregnancy.

Use in lactation:

- Excreted in breast milk.
- Maternal ingestion of paracetamol in normal therapeutic doses does not appear to present a risk to the nursing infant.

Adverse Effects / Undesirable Effects: Adverse effects of paracetamol are rare and usually mild, although haematological reactions have been reported. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Overdose and Treatment

Symptoms: Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. The most serious adverse effect of acute overdose of paracetamol is a dose-dependent, potentially fatal hepatic necrosis. Clinical and laboratory evidence of hepatotoxicity may be delayed for up to one week. Major manifestations of liver failure such as jaundice, hypoglycemia and metabolic acidosis may take at least 3 days to develop.

Treatment: In cases of overdose, methods of reducing the absorption of ingested drug are important. Gastric lavage is essential even if several hours have elapsed. Prompt administration of 50g activated charcoal and 500ml iced mannitol 20% by mouth, may reduce absorption. If the history suggests that 15g Paracetamol or more has been ingested, administer one of the following antidotes:

Acetylcysteine 20% i.v.: Administer intravenously, 20% acetylcysteine immediately without waiting for positive urine test or plasma level results: initial dose of 150mg/kg over 15 minutes, followed by continuous infusion of 50mg/kg in 500ml 5% glucose/dextrose over 4 hours and 100mg/kg in 1L 5% glucose/dextrose over 16 hours; OR

Oral Methionine: 2.5g immediately followed by three further doses of 2.5g at four hourly intervals; OR

Oral Acetylcysteine 5%: 140mg/kg as a loading dose, then 70mg/kg every 4 hours for a total of 17 maintenance doses. If more than ten hours have elapsed since the overdosage was taken, the antidote may be ineffective.

Storage Conditions: Store below 30°C.

Shelf-Life: Three years from the date of manufacture.

Dosage Forms and Packaging Available: Blister pack of 10's per strip. Box of Box of 3x10's and 20x10's

Manufacturer & product registration holder

AV MANUFACTURING SDN BHD (667760-V)
Lot 10621 (PT16700), Jalan Permata 2, Arab Malaysian Industrial Park, 71800 Nilai, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

Jun 2024 Ver.03

Risalah Maklumat Ubat untuk Pengguna (RiMUP) - Avadol®-Fast Tablet 500mg (Bahan aktif: Paracetamol 500mg)

1. Apakah kegunaan Avadol-Fast: Avadol-Fast digunakan untuk melegakan demam, kesakitan dari ringan kepada sederhana termasuk: sakit kepala, migrain, sakit belakang, sakit otot , myalgia, senggugut, sakit osteoarthritis, sakit gigi, ketidakselesaan dari selsema, dan sakit tekak.

2. Bagaimana Avadol-Fast Berfungsi: Avadol-Fast diserap dengan cepat melalui pengambilan dari mulut. Ia diserap dalam usus kecil. Paracetamol juga dikumuh dengan cepat melalui urin.

3. Sebelum mengambil Avadol-Fast

Bila tidak boleh mengambil

Jika anda

- hipersensitiviti terhadap paracetamol atau bahan-bahan yang terkandung dalam produk ini.
- sedang mengambil ubat yang lain juga mengandungi paracetamol.
- menghadapi kemerosotan fungsi hati yang teruk.

Sebelum mula mengambil

Semak dengan doktor anda, jika anda

- menghadapi sebarang penyakit ginjal atau hati, termasuk penyakit hati alkohol.
- menghadapi artritis yang sederhana dan perlu mengambil ubat tahan sakit setiap hari.
- mempunyai buah pinggang atau hati yang lemah.

Jika mengambil ubat-ubat lain

Sila beritahu doktor atau ahli farmasi jika anda ada mengambil ubat-ubatan lain seperti

- Anti-pembeku (seperti warfarin) dan anda perlu mengambil ubat tahan sakit setiap hari.
- Ubat-ubatan lain yang anda sedang guna, seperti ubat dibeli tanpa preskripsi.

Hamil dan penyusuan anak

Penggunaan semasa hamil:

- Dianggap pilihan analgesik terhadap pesakit yang hamil.
- Walaupun ia akan melalui plasenta, paracetamol dianggap selamat dalam normal dos terapeutik untuk tempoh yang pendek sebagai minor analgesic/antipiretik dalam kehamilan.

Kegunaan dalam penyusuan:

- Didapati dalam susu ibu jika ibu menggunakan.
- Pengambilan paracetamol mengikut dos terapeutik oleh ibu, tidak akan membawa risiko kepada bayi yang diberi susu.

Memandu kereta dan mengendali mesin

Paracetamol mempunyai potensi yang kecil untuk menjejaskan keupayaan seseorang pesakit mengendali mesin atau memandu.

4. Cara mengambil Avadol-Fast

Berapa banyak harus digunakan

Dewasa: Ambil 1-2 biji untuk 3 atau 4 kali sehari. Maksimum: 8 biji sehari. Tidak sesuai untuk kanak-kanak.

Bagaimana boleh diambil: Avadol-fast tablet patut ditelan dengan air.

Berapa lama perlu diambil: Tempoh rawatan ditentukan oleh doktor anda.

Jika terlupa mengambil: Ambil dos yang seterusnya. Jangan mengambil dos berganda untuk menganti dos yang terlupa ambil.

Jika mengambil berlebihan: Hubungi doktor anda dengan segera atau pergi ke Jabatan Kecemasan hospital terdekat jika terambil dos berlebihan. Ini perlu dilakukan walaupun tiada sebarang kesan keracunan atau rasa tidak selesa. Anda mungkin memerlukan perhatian perubatan segera.

5. Semasa mengambil Avadol-Fast

Perkara yang perlu dilakukan: Simpan RiMUP, anda mungkin perlu membaca semula. Jika anda ada sebarang pertanyaan, sila tanya doktor atau ahli farmasi.

Perkara yang tidak boleh dilakukan: Ubat ini adalah preskripsi untuk anda sahaja. Jangan bagi orang lain. Ia mungkin tidak sesuai untuk orang lain walaupun mereka mempunyai simpton-simpton yang sama.

Perkara yang perlu diberi perhatian: Jangan mengambil melebihi dos yang disyorkan. Jika simptom-simptom kekal (jika kesakitan tersebut mengekal 10 hari ke atas) atau sakit kepala kekal, sila jumpa doktor anda.

6. Kesan-kesan Sampingan: Kesan-kesan sampingan paracetamol adalah jarang dan biasanya ringan, walaupun reaksi hematologic pernah dilaporkan. Tindak balas hypersensitiviti kulit termasuk ruam kulit, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis telah dilaporkan.

Amaran alahan: Paracetamol mungkin menyebabkan reaksi kulit yang serius. Simptom-simptom termasuk kemerahan kulit, lepuh dan ruam. Ini merupakan simpton-simpton keadaan yang serius. Jika berlaku tanda-tanda tersebut, sila berhenti dan mendapatkan rawatan segera.

Anda boleh melaporkan sebarang kesan sampingan atau kesan advers ubat kepada Pusat Pemantauan

Kesan Advers Ubat Kebangsaan melalui laman web npra.gov.my

[Consumer—>Reporting Side Effects to Medicines (ConSERF)or Vaccines (AEFI)]

7. Cara Penyimpanan dan Pelupusan: Simpan bawah 30°C. Jangan menggunakan jika tarikh yang dicetak pada komponen pembungkusan telah luput. Ubat tidak boleh dilupuskan melalui air buangan atau buangan isi rumah. Sila bertanya kepada ahli farmasi anda bagaimana untuk melupuskan ubat yang tidak lagi diperlukan. Langkah-langkah ini akan membantu melindungi persekitaran.

Jauhi dari kanak-kanak

8. Maklumat Lanjut

Rupa and wama produk: Avadol-Fast tablet berbentuk bulat, berwarna putih dengan 500 pada satu permukaan, dan licin pada permukaan belakang.

Bahan-bahan kandungan

Bahan Aktif: Paraceramol 500mg

Nombor Pendaftaran: Avadol-Fast Tablet 500mg (MAL15125042XZ)

9. Pengilang & pemegang pendaftaran produk

AV Manufacturing Sdn. Bhd. (667760-v)

Lot 10621 (PT16700), Jalan Permata 2, Arab Malaysian Industrial Park, 71800 Nilai, Negeri Sembilan, Malaysia.

Nombor siri : NPRA/R(2/1) 220726/923

Ubat ini mengandungi PARACETAMOL.
Jangan mengambil ubat lain yang mengandungi Paracetamol pada masa yang sama.

此药物含有PARACETAMOL.
请不要同时服用其他拥有Paracetamol的药物。

消费者药物信息 (RiMUP) - Avadol®-Fast Tablet 500mg (有效成份: Paracetamol 500 mg)

1.Avadol-Fast的用途: Avadol-Fast有助于缓解发烧，轻度至中度疼痛，包括：头痛，偏头痛，背痛，肌肉骨骼疼痛，肌痛，痛经（周期性疼痛），骨关节炎疼痛，牙痛以及感冒，流感和咳嗽痛。

2.Avadol-Fast如何运作:Avadol-Fast通过口服迅速被吸收，吸收主要发生在小肠中。Paracetamol的排泄迅速且通过尿液发生。

3.在使用Avadol-Fast之前

你不能使用它如果你曾有过

- 对Avadol-Fast的任何其他成份过敏。
- 服用含paracetamol的其他药物。

•严重和积极的肝损害。

在你开始使用它之前

与您的医生核对

- 如果您患有肝脏或肾脏问题，包括酒精性肝病
- 如果您患有轻微关节炎并且需要每天服用止痛药
- 如果您的肾脏或肝功能受损。

服用另一种药物

如果您正在服用这些药物，请咨询您的医生或药剂师：

- 如果您正在服用血液稀释药物（抗凝剂，如华法林），您需要每天服用止痛药
- 如果您最近服用了其他药物，包括未经处方获得的药物。

怀孕和哺乳

怀孕期间使用：

- 它被认为是孕妇可以使用的止痛药。
- 尽管扑热息痛可以穿过胎盘，但短时间与正常剂量服用作为止痛药或发烧药是安全的。

喂食期间的用途：

- 母亲服用的扑热息痛将存在于母乳中，但常规剂量不会伤害婴儿。

驾驶和使用机器:这不可能损害患者驾驶或使用机器的能力。

4.如何使用Avadol-Fast

使用量:成人：每天服用1至2片，每次3至4次。最多：每天8片。不建议儿童使用。

如何使用它:Avadol-Fast片剂应该用水吞服。

使用它多久:Avadol-Fast治疗的持续时间由您的医生决定。

如果你忘记使用它:请直接服用下一剂。不要服用双倍剂量来弥补忘记的剂量。

如果您使用过量:请立即到邻近的医院就医。

5.当你使用Avadol-Fast时

你必须这样做:保留此传单。你可能需要把它重读一遍。如果您还有其他问题，请咨询您的医生或药剂师

不能做的事情:这种药已经为你处方了。不要将它传递给其他人。它可能伤害他们，即使他们的症状与你的相同。

要小心事情:不要超过推荐剂量。如果您的症状持续存在（如果疼痛持续10天以上）或您的头痛持续存在，请与您的医生见面。

6.副作用:Paracetamol的副作用很少见，通常很轻微，尽管血液学反应已有报道。

皮肤过敏反应包括皮疹，血管性水肿，史蒂文斯约翰逊综合征中毒性表皮坏死已经有报道

过敏警报: Paracetamol可能导致严重的皮肤反应。症状可能包括皮肤发红，水泡或皮疹。

这些可能是严重情况的迹象。如果发生这些反应，请立即停止使用并寻求医疗援助。

您可以通过浏览网页 npra.gov.my [Consumers—>Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]，直接向国家药物不良反应检测中心报告任何副作用或不良反应。

7.储存和处置

存放在30°C以下。请勿在包装组件上印刷的有效日期之后使用。

不应该通过废水或生活垃圾处理药物。询问药剂师如何处理不再需要的药物。这些措施将有助于保护环境。

请将本品放在儿童不能接触的地方。

8.产品说明

它看起来像什么: Avadol-Fast是白色，圆形，一面标有“500”字样，另一面平滑。

成份

有效成份: Paracetamol 500mg

注册号码: Avadol-Fast Tablet 500mg (MAL15125042XZ)

9. 制造商和产品注册持有人

AV MANUFACTURING SDN BHD (667760-v)

Lot 10621 (PT16700), Jalan Permata 2, Arab Malaysian Industrial Park, 71800 Nilai, Negeri Sembilan, Malaysia.

Nombor siri : NPRA/R(2/1) 220726/923