

QUEMOX POWDER

50MG/G

Ingredient(s)

Each gram contains:

Mebendazole.....50mg

Sodium benzoate (as preservative).....1mg

Pharmacodynamics

Mebendazole is an anthelmintic belonging to the benzimidazole (BZD) group of compounds with efficacy against certain gastrointestinal roundworms and lungworms. The BZD act primarily by binding to nematode tubulin. Specifically, BZD bind to beta-tubulin, which in turn prevents its dimerization with alpha-tubulin and the polymerization of tubulin oligomers into microtubules. Microtubules are essential structure units of many organelles and are necessary for numerous cellular process, including mitosis, protein assembly, and energy metabolism. Mebendazole acts selectively against the gastrointestinal and lung parasites in the host, leading to an irreversible lytic necrosis of those cells and death of the worm. Efficacy of mebendazole (30ppm in feed for 10 days) expels greater than 99% of swine whipworms. Mebendazole has demonstrated positive results against larval stage of *T. hydatigena* in pigs. A brief regimen of treatment (25mg/kg bw/day) for 5 days was totally effective for Taenia cyclicerci in pigs.

Pharmacokinetics

Mebendazole is poorly metabolized and most is excreted unchanged in faeces within 24-48hours. Between 5 and 10% is excreted in urine and only a small amount is excreted as the decarboxylated derivative of mebendazole. The pharmacokinetic profile of mebendazole is similar in various animal species, including pigs. Mebendazole has a poor oral bioavailability, due to a low solubility in aqueous systems, a slow dissolution rate in the gastrointestinal tract and first-pass metabolism in the gut wall and the liver. This causes a high faecal excretion of parent drug and low levels in plasma and tissues.

Indications

An oral broad spectrum anthelmintic for the treatment of helminthiasis in pig. The product is effective against gastrointestinal worms: *Ascaris suum* larvae including *Oesophagostomum dentatum*, *Hyoststrongylus rubidus*, *Trichuris suis*, *Strongyloides ransomi* , *Globocephalus sp*; and lung worms : *Metastrongylus*

Target Species

Pig (Piglets, fattening pigs, sows, boars).

Recommended Dose

- 1.Piglets, fattening pigs: Mix 60g of Quemox Powder to 100kg of complete feed (30 ppm). Give for 5 days (in severe *Strongyloides* or *Trichuris infection*: 10 days).
- 2.Sows: Mix 60g of Quemox Powder to 100kg of complete feed (30 ppm). Give for 10 days.
- 3.Boars: 0.3 g of Quemox Powder per 10 kg body weight daily for two consecutive days in the feed mix.

Mode of Administration

Orally. Add to feed.

Contraindication

Do not use in animals with hypersensitivity to the active ingredient.

Warning and Precautions

Care should be taken to avoid the following practices due to increased risk of development of resistance and could ultimately result in ineffective therapy: Too frequent and repeated use of anthelmintics from the same class, over an extended period of time. Underdosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device (if any). Suspected clinical cases of resistance to anthelmintics should be further investigated. Resistance to mebendazole (a benzimidazole) has been reported in cyathostomes in horses. Therefore, the use of this product should be based on local epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

i. Special precautions for use in animals

Not known.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use

Persons with known hypersensitivity to ingredients in the product should not handle the product. Avoid contact with skin and eyes. Wash off any contact with the product immediately with clean running water. Seek medical attention if irritation persists.

Interaction with Other Medicaments

Not known.

Pregnancy and Lactation

Not known.

Side Effects

Pigs ,well tolerated. Taking more than 30 ppm may cause temporarily softening of the faeces.

Environmental Property

Any unused product or waste material should be disposed in accordance with national requirements.

Symptoms and Treatment of Overdose

Supportive therapy if required.

Storage Condition

Store at temperature below 30°C. Protect from direct sunlight and moisture.

Shelf Life

4 years

Withdrawal Periods

2 days.

Maximum Residual Limit (MRL)

Pig : liver: 0.02 mg/kg.

Product Description

A white to off-white color powder, and without other foreign matter.

600g aluminum pouch.

20kg paper bag.



Made by YSP

Manufacturer:

Y.S.P. INDUSTRIES VIETNAM CO., LTD

Road No. 6, Nhon Trach 3 Industrial Zone, Nhon Trach District,
Dong Nai Province , Vietnam. Tel: +8461 3569 993

Product Registration Holder :

Y.S.P. INDUSTRIES (M) SDN. BHD.

Lot 3, 5 & 7, Jalan P/7, Section 13,

Kawasan Perindustrian Bandar Baru Bangi,

43000 Kajang, Selangor Darul Ehsan Malaysia.

Ordering Line: 1 800 88 3027 Product Info Line: 1 800 88 3679