

hovid-Paracetamol Syrup 120mg / 5mL (SUGAR FREE)

VIPARXX-P4

DESCRIPTION

Orange liquid with orange flavor.

COMPOSITION

Each 5 mL contains: Paracetamol 120mg

ACTIONS AND PHARMACOLOGY

Paracetamol is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties.

- **Analgesic:** The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (specifically cyclooxygenase (COX)-2) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation.

The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation.

- **Antipyretic:** Paracetamol act centrally on the hypothalamic heat-regulating center to produce peripheral vasodilatation resulting in increase blood flow through the skin, sweating and heat loss. Paracetamol reduces fever by inhibiting the formulation and release of prostaglandins in the CNS and by inhibiting endogenous pyrogens at the hypothalamic thermoregulator center.

PHARMACOKINETICS

Following oral administration paracetamol is rapidly absorbed.

Paracetamol absorption takes place mainly in the small intestine and therefore the rate of absorption is depending on the rate of gastric emptying. It has been shown that drugs which delay gastric emptying also delay the absorption of paracetamol whereas metoclopramide (a drug which increases the rate of gastric emptying) accelerates absorption of the analgesic through the total amount absorbed doses not increase.

The presence of food in the stomach has also been reported to reduce the rate of absorption of paracetamol. Alterations in gastric pH have no appreciable effect on paracetamol absorption. During absorption, the amount of paracetamol which is inactivated is negligible and it has been shown that paracetamol dose not affect gastric mucosal permeability and does not produce mucosal bleeding.

Peak plasma concentrations are reached 1 hour after absorption. The plasma half life is 1 to 3 hours.

Paracetamol penetrates the brain and is present in breast milk of human.

Paracetamol is metabolized by the microsomal enzyme system of the liver. This metabolism is mainly to the glucuronide and sulphate conjugates, accounting for approximately 49% and 26% of the ingested dose respectively. About 4% is excreted as free paracetamol. Other minor pathways include the production of catechol derivatives and cysteine conjugates (via glutathione). Paracetamol excretion is rapid and occurs via the urine.

INDICATION

Indicated for the relief of fever, headache and symptoms of cold and flu, toothache, discomfort of teething and fever after vaccination.

CONTRAINDICATION

- Hypersensitivity to paracetamol or any of the other ingredients/ components of the product
- Severe and active hepatic impairment

WARNINGS AND PRECAUTIONS

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time

- Keep out of reach of children.
- Do not take if allergic to paracetamol.
- Patients should contact their health care provider if symptoms persist (if the pain lasts for more than 10 days, if there is redness or fever lasts more than 3 days).
- Paracetamol should be given with care to patients with impaired kidney or liver function.
- Large doses should be avoided in patients with hepatic impairment. Paracetamol overdose may harm the liver.
- Do not exceed recommended dose.
- Paracetamol provides symptomatic relief only, additional therapy to treat the cause of the pain or fever should be instituted when necessary.
- This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.
- **Allergy alert:**
Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.
These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Effects on Ability to Drive and Use Machines

It is unlikely to impair a patient's ability to drive or use machinery.

PREGNANCY AND LACTATION

Pregnancy

- Considered to be the analgesic of choice in pregnant patients.
- Although it crosses placenta, paracetamol is considered to be safe in normal therapeutic doses for short-term use as a minor analgesic/antipyretic in pregnancy.

Lactation

- Excreted in breast milk.
- Maternal ingestion of paracetamol in normal therapeutic doses does not appear to present a risk to the nursing infant.

DRUGS INTERACTIONS

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

MAIN SIDE/ ADVERSE EFFECTS

Adverse effects of paracetamol are rare and usually mild, although haematological reactions have been reported. Skin rashes and other hypersensitivity reactions occur occasionally.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

OVERDOSE AND TREATMENT**Symptoms:**

Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. The most serious adverse effect of acute overdose of paracetamol is a dose-dependent, potentially fatal hepatic necrosis. Clinical and laboratory evidence of hepatotoxicity may be delayed for up to one week.

Major manifestations of liver failure such as jaundice, hypoglycemia and metabolic acidosis may take at least 3 days to develop.

Treatment:

In cases of overdose, methods of reducing the absorption of ingested drug are important. Gastric lavage is essential even if several hours have elapsed. Prompt administration of 50g activated charcoal and 500ml iced mannitol 20% by mouth, may reduce absorption.

If the history suggests that 15g Paracetamol or more has been ingested, administer one of the following antidotes:

- **Acetylcysteine 20% i.v.:** Administer intravenously, 20% acetylcysteine (Parvolex, Glaxo) immediately without waiting for positive urine test or plasma level results: initial dose of 150mg/kg over 15 minutes, followed by continuous infusion of 50mg/kg in 500ml 5% glucose/dextrose over 4 hours and 100mg/kg in 1L 5% glucose/dextrose over 16 hours; OR
- **Oral Methionine:** 2.5g immediately followed by three further doses of 2.5g at four hourly intervals. For a 3 year old child, 1g methionine every four hours for four doses has been used; OR
- **Oral Acetylcysteine 5%:** 140mg/kg as a loading dose, then 70mg/kg every 4 hours for a total of 17 maintenance doses. If more than ten hours have elapsed since the overdosage was taken, the antidote may be ineffective.

DOSAGE AND ADMINISTRATION

Oral.

hovid-Paracetamol Syrup (Sugar Free) may be taken with or without food.

hovid-Paracetamol Syrup 120mg / 5mL (Sugar Free)

For children aged 1 to 6 years. Give in 6 hourly intervals; not more than 4 times a day.

AGE	PARACETAMOL SYRUP 120mg /5mL
1 – 3 years	6ml – 10ml
4 – 6 years	10ml – 15ml

The above table is based on the average weight-age grid of children. If you know the body weight of your child, it's always more accurate to use the weight to arrive at the correct dose. Please refer below.

Using the body weight of your child, you can calculate the dose of Paracetamol Syrup 120mg/5ml your child needs. Calculation is based on recommended dose of 15mg per body weight in kg (BW).

Table below provides the dose of Paracetamol Syrup 120mg/5ml to be given, based on body weight:

CHILD'S WEIGHT IN KG	IN IB	PARACETAMOL SYRUP 120mg /5mL
10	22	6 mL
15	33	9 mL
20	44	13 mL
30	66	19 mL

Note: 1kg = 2.2lb

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C. Do not refrigerate.

Presentation/ Packing:

Syrup 120mg / 5ml x 60ml, 100ml and 120ml.

Product Registration Holder / Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Malaysia.

Information date: December 2018