

SG.OXYTETRA LA 20% (w/v) Solution

For Animal Use Only

Name and Strength of Active Substance(s)

Oxytetracycline Dihydrate 216 mg/ml (Equivalent to Oxytetracycline 200 mg/ml)

Product Description:

SG.Oxytetra LA 20% (w/v) Solution is sterile, clear, stable, ready-to-use solution. It is a solution with brown to dark brown color. It is consistency solution, stability at pH from 8 to 9.

Pharmacodynamics:

Oxytetracycline is a bacteriostatic antibiotic that inhibits protein synthesis in susceptible bacteria. Inside the cell it binds irreversibly to receptors on the 30S subunit of the bacterial ribosome where it interferes with the binding of the aminoacyl-transfer RNA to the acceptor site on the messenger RNA ribosome complex. This effectively prevents the addition of amino acids to the elongating peptide chain, inhibiting protein synthesis. The product is specifically formulated to provide a prolonged action resulting in sustained antibacterial activity.

Oxytetracycline has been shown to be effective in vitro against the following bacterial species: *Bordetella bronchiseptica*, *Corynebacterium pyogenes*, *Erysipelothrix rhusiopathiae*, *Escherichia coli*, *Haemophilus somnus*, *Pasteurella haemolytica*, *Pasteurella multocida*, *Salmonella dublin*, *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus faecalis*, *Streptococcus pyogenes* and *Streptococcus uberis*.

Pharmacokinetics:

Blood levels persist for at least 4 days after administration by the intramuscular route. Maximum blood levels are achieved between 4 and 8 hours following intramuscular administration.

Indications:

The product is indicated for use in cattle, sheep and pigs in the treatment of:

- Atrophic rhinitis caused by *Bordetella bronchiseptica*, *Mannheimia haemolytica* and *Pasteurella multocida*.
- Navel/joint ill caused by *Arcanobacterium pyogenes*, *E. coli* or *Staphylococcus aureus*.
- Mastitis caused by *Corynebacterium pyogenes*, *E. coli*, *Staphylococcus aureus*, *Streptococcus agalactiae* or *Streptococcus uberis*.
- Metritis caused by *E. coli* or *Streptococcus pyogenes*.
- Pasteurellosis and infections of the respiratory tract caused by *Mannheimia haemolytica* and *Pasteurella multocida*.
- Septicaemia caused by *Salmonella dublin* and *Streptococcus pyogenes*.
- Erysipelas caused by *Erysipelothrix rhusiopathiae*.

The product can also be used in the control of enzootic abortion in sheep.

Recommended Dosage:

The recommended dose rate is 20 mg/kg bodyweight (i.e. 1 ml per 10 kg body weight) administered by deep intramuscular injection. The product is recommended for a single administration only. Maximum recommended dose at any one site:

Cattle:	20 ml
Pigs:	10 ml
Sheep:	5 ml
Piglets:	1 day 0.2 ml
	7 days 0.3 ml
	14 days 0.4 ml
	21 days 0.5 ml
	Over 21 days 1.0 ml/10 kg.

Mode of Administration:

Intramuscular injection, single administration only.

Contraindication:

- Not for use in horses, dogs, cats.
- Not for use in animals suffering from hepatic or renal damage.

Warning and Precautions:

Special precautions for use in animals

- Do not dilute. If concurrent treatment is administered use a separate injection site.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

- Wash hands after use.
- In case of contact with eyes or skin, wash immediately with water as irritation may occur.
- Take care to avoid accidental injection.

Interaction with Other Medicinal Product: None known.

Pregnancy and Lactation:

The use of oxytetracycline during the period of tooth and bone development, including late pregnancy may lead to tooth discoloration, the product can be safely administered to lactating animals.

Adverse Effects and Toxicity:

Although the product is well tolerated, occasionally a slight local reaction of a transient nature has been observed.

Overdose and Treatment:

There is no known specific antidote, if signs of possible overdose occur, treat the animal symptomatically.

Withdrawal Period:Cattle

Meat – 31 days, Milk – 10 days

Sheep

Meat – 9 days, Milk – 7 days

Pigs

Meat – 18 days

Incompatibilities: None known.

Storage Condition:

Store in a dry place at room temperature (below 30 °C), keep away from direct sunlight. Keep out of reach of children (*Jauhkan daripada kanak-kanak*).

Proposed shelf life (first opening of container):

28 days

Dosage Form:

Solution

Packaging Available:

100ml

Date of Revision of Package Insert: 15th April 2025

Name & Address of Manufacturer:

SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY
Anova Corporation Industrial zone, Long Cang commune, Tay Ninh province, Vietnam.

Changed the manufacturer name and update address from:

ANOVA PHARMA JOINT STOCK COMPANY

Anova Corporation Industrial Zone, Long Cang Commune, Can Duoc District, Long An Province, Vietnam- Tel:(+84 272)3726262.

to:

SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY

Anova Corporation Industrial zone, Long Cang commune, Tay Ninh province, Vietnam- Tel:(+84 272)3726262.

Marketing Authorisation Holder:

SUNZEN CORPORATION SDN BHD

No. 11, Jalan Anggerik Mokara 31/47, Kota Kemuning, 40460 Shah Alam, Selangor, MALAYSIA.