

### **D3.0 Proposed package insert**

#### **Stalleng Nystatin Premix Powder**

**123.2miu/kg**  
**(FOR ANIMAL USE ONLY)**  
Anti-fungal Feed Additive

#### **DESCRIPTION**

Stalleng Nystatin Premix Powder 123.2miu/kg is a free flowing, greyish powder with slight characteristic odour of fermentation from the active ingredient Nystatin

#### **COMPOSITION/KG:-**

Nystatin .....123.2miu  
Calcium Carbonate..... 956 gm

#### **INDICATIONS**

Use as an aid in control or treatment of Crop Mycosis, Mycotic Diarrhoea (Candida albicans) in Turkey

#### **PHARMACODYNAMICS**

Nystatin has a mechanism of action similar to that of amphotericin B. It binds to sterols in the membrane of the fungal cell altering the permeability of the membrane allowing intracellular potassium and other cellular constituents to "leak out". When given orally, the drug must come into contact with the organism to be effective. Nystatin has activity against a variety of fungal organisms, but is clinically used against topical, oropharyngeal, and gastrointestinal Candida infections

#### **PHARMACOKINETIC**

Nystatin is not measurably absorbed after oral administration and almost entirely excreted unchanged in the faeces. The drug is not used parentally because it is reportedly extremely toxic to internal tissues.

#### **CONTRA INDICATION/ Precaution & Warning**

Nystatin is contraindicated in animals with known hypersensitivity to it. Do not confuse Nystatin with HMG-CoA reductase inhibitors ("statin drugs", e.g., simvastatin, etc.)

#### **SIDE EFFECTS**

Occasionally, high dosages of nystatin may cause GI upset (anorexia, vomiting, and diarrhea). If suspected adverse reaction occur, treatment should be discontinued and to inform veterinarian/local authority.

#### **PREGNANCY AND LACTATION**

Although the safety of the drug during pregnancy has not been firmly established, the lack of appreciable absorption or case reports associating the drug with teratogenic effects appears to make it safe to use.

#### **SYMPTOMS AND TREATMENT OF OVERDOSE**

Because the drug is not absorbed after oral administration, acute toxicity after an oral overdose is extremely unlikely, but transient GI distress may result.

#### **INTERACTION WITH OTHER MEDICAMENTS**

No significant interactions reported for oral Nystatin.

**STORAGE:** Store in cool dry place at 30°C and below, away from direct sunlight

**PROPOSED SHELF LIFE (as packaged for sale)**  
2 years (24 months)

**PROPOSED SHELF LIFE (AFTER FIRST OPENING OF CONTAINER)**  
24 hours

#### **ADMINISTRATION:-**

Oral route by mixing the powder into animal feed ration.

#### **DOSAGES:**

Use as an aid in control of crop mycosis and mycotic diarrhoea (Candida albicans):  
1.136kg (equivalent to 140miu or 50g of Nystatin) per ton of feed.

Treatment of crop mycosis and mycotic diarrhoea (Candida albicans):  
2.273kg (equivalent to 280miu or 100g of Nystatin) per ton of feed for 7 to 10 days."

**KEEP OUT OF REACH OF CHILDREN**  
**JAUHKAN UBAT DARI KANAK-KANAK**

#### **WITHDRAWAL PERIOD:**

No withdrawal period is required.

**PACKING:** 25kg multilayer with inner LDPE layered paper bag.

#### **MANUFACTURED BY:**

Range Pharma Sdn Bhd, (144401U)  
No.1, Jalan TSB 11, Taman Industri Sungai Buloh, 47000 Sungai Buloh. Malaysia.

#### **REGISTERED AND MARKETED BY:**

Stalleng (M) Sdn Bhd (408941-X),  
23, Jalan Serindit 2,  
Puchong Jaya Industrial Park,  
47100 Puchong, Selangor.  
Telephone: 03-58822968 Fax: 03-58826778

Date of Revision: 06/04/2017