

PELANGI VET PHARMA SDN.BHD (665097-K)
LOT 8, PERDA Industrial Park, Lorong IKS, Simpang Ampat B,
14100 Simpang Ampat, S.P.S, Pulau Pinang, Malaysia

PRODUCT LITERATURE

1. Product Name:

PVP- CTC 20% W/W POWDER

2. Each 1g powder contains:

Active Ingredient:

Chlortetracycline Hydrochloride BP 200.00mg

3. Description

Yellowish powder

4. Target Species

Pigs

5. Pharmacodynamics:

Chlortetracycline hydrochloride is a predominantly bacteriostatic antibiotic, interfering with bacterial protein synthesis of the rapidly growing and reproducing bacterial cell. Chlortetracycline has a broad spectrum of activity, including Gram-positive aerobes, Gram-negative anaerobes and Mycoplasmas. Resistance is known to occur in respiratory pathogens of pigs and cross-resistance occurs between chlortetracycline and other tetracyclines.

6. Pharmacokinetics:

Following oral administration, maximum blood levels are achieved within approximately 2-8 hours. Steady state plasma concentrations of chlortetracycline are maintained throughout the twice daily seven day treatment period.

7. Indication:

Pigs:

The product is indicated in the treatment of respiratory disease in pigs caused by micro-organisms sensitive to chlortetracycline.

8. Recommended Dose:

For oral administration.

The recommended therapeutic dose is 20 mg chlortetracycline hydrochloride per kg bodyweight (equivalent to 15 grams of PVP-CTC 20% w/w powder per 150 kg bodyweight) per day administered for seven days.

This should be given in a divided daily dose i.e. 7.5 g morning and 7.5 g evening. As a guide to dosing pigs of different weight, see table.

The product should be administered to small quantities of feed for immediate consumption by individual animals. Larger groups should be treated with medicated feeding stuff.

The product should be mixed thoroughly into a part of the daily feed ration and should be administered prior to the feeding. It should be ensured that the calculated dose is completely taken up by the animal.

If animals don't recover within 3 days after oral medication, diagnosis should be reconsidered and treatment should be changed, if necessary.

To ensure the correct dosage and to avoid possible under-dosing, the bodyweight and the amount of product administered should be determined as accurately as possible. To determine the correct amount of product, a calibrated weighing scale should be used.

Pig Bodyweight (kg)	Daily amount (g) of PVP-CTC 20% w/w Powder	Dose (g) – to be given TWICE daily
15 kg	1.5g	0.75g
30 kg	3g	1.5g
60 kg	6g	3g
75 kg	7.5g	3.75g
150 kg	15g	7.5g

9. Mode of Administration:

Oral

10. Contraindications:

Do not use in animals with known hypersensitivity to tetracyclines or any of the excipients.

Do not use in animals with severe liver and renal disorders.

11. Warning & Precautions:**Special warnings for each target species**

The uptake of oral medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed, animals should be treated parenterally.

Special precautions for use**Special precautions for use in animals**

The product is efficient only against bacterial strains most sensitive to chlortetracycline. Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Inappropriate use of the product may increase the prevalence of bacteria resistant to chlortetracycline and may decrease the effectiveness of treatment with related substances, due to the potential for cross-resistance.

Long term use of this product is not recommended as it may lead to the development of bacterial resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Handle this product with care to avoid exposure when adding to feed and administering medicated feed to the animals.

Take adequate measures to avoid dust formation when adding the product to feed.

Those handling the product should do so in a mechanically ventilated area.

Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143.

Direct contact of the product with the skin, eyes and mucous membranes should be avoided.

Wear protective gloves, overalls and approved safety glasses. In case of accidental exposure, wash area immediately with water.

Do not smoke, eat or drink when handling the product.

Hands and exposed skin should be washed thoroughly after use.

Other precautions

None.

12. Drug Interactions:

This product is not recommended for concurrent administration with any other oral medication.

Do not add the product to feed overloaded with polyvalent cations such as Ca^{2+} and Fe^{3+} because the formation of chlortetracycline complexes with these cations is possible. Do not administer together with antacids, kaolin and iron preparations and in conjunction with bactericidal antibiotics like beta-lactams.

13. Pregnancy & Lactation:

The use is not recommended during pregnancy or lactation.

The treatment of pregnant animals with chlortetracycline may result in adverse effects on skeletal and tooth development in the foetus. Therefore, the product should be used only in pregnant sows according to the benefit/risk assessment of the responsible veterinarian.

14. Side effects:

Chlortetracycline toxicity is low. If digestive disturbances do occur, treatment should be discontinued.

On rare occasions (more than 1 but less than 10 animals in 10,000 animals) the following adverse reactions may occur: allergic reactions and photosensitivity; gastrointestinal disorders; disorders of the liver and the kidneys. If suspected adverse reactions occur, treatment should be discontinued.

15. Symptoms & Treatment of Overdose:

Do not exceed the stated dose.

Chlortetracycline toxicity is low. If digestive disturbances do occur, treatment should be discontinued.

16. Withdrawal Period:

Pigs:

Meat and offal: 6 days

17. Storage Conditions:

Store below 30° C in dry place, protected from light.

18. Dosage Form:

Powder in the form of powder to be mixed in feed.

19. Packaging:

- (i) Pack of 1kg in HDPE Plastic Container along with caps.
- (ii) Pack of 500g in HDPE Plastic Container along with caps.
- (iii) Pack of 1kg in Aluminium Sheet Lacquered .
- (iv) Pack of 500g in Aluminium Sheet Lacquered.

20. Disposal of Container:

To be disposed as per local authority requirement.

21. Expiry date:

2 years from date of manufacturing

22. Shelf-life after reconstitution or dilution:

To be used within 24 hours after dilution.

Manufactured & Distributed by:

Pelangi Vet Pharma Sdn Bhd. (665097-K)

Lot 8, PERDA Industrial Park,

Lorong IKS, SimpangAmpat B

14100 SimpangAmpat,

S.P.S Pulau Pinang, Malaysia

Last update on: 15 March 2017