

PRODUCT LITERATURE

1. Product Name:

PVP- DIAPRIM SOLUBLE POWDER

2. Each 1g powder contains:

Active Ingredient:

Sulfadiazine/Sulphadiazine BP 400.00mg

Trimethoprim BP 80.00mg

3. Description

A white to off white powder

4. Target Species

Chickens, turkeys and pigeons.

5. Pharmacodynamics:

Mechanism of Action:

Trimethoprim and sulfadiazine are antimicrobial substances which act sequentially in the synthesis of tetrahydrofolic acid from para-aminobenzoic acid. The action of trimethoprim is to potentiate the sulfadiazine component. Veterinary preparations contain the two active substances in a 1:5 ratio and this combination is generally effective clinically because of the relatively broad range of drug ratio over which synergism occurs

6. Pharmacokinetics:

Both active substances are absorbed well and distributed throughout the body. Excretion occurs mainly via the urine.

7. Indication:

For the control and treatment of sensitive bacteria involved in yolk sac and navel infections, pericarditis, clinical salmonella infections; Fowl cholera (*Pasteurella multocida*) and *Escherichia coli* problems which arise as complications in respiratory disease of chickens, turkeys and pigeons.

8. Recommended Dose:

CHICKEN AND TURKEYS:

Birds less than 14 days; use dose rate of 1g per 16kg bodyweight daily for 3 to 5 days.

Birds over 14 days and less than 1kg, use dose rate of 1g per 32 kg bodyweight daily for 3 to 5 days.

Birds greater than 1kg, use dose rate of 1g per 16kg bodyweight for the first 24 hours of treatment. For each day of treatment thereafter, administer at 1g per 32kg bodyweight each day for the remaining 3 to 4 days.

PIGEONS:

For adult birds, use dose rate of 1g per 16kg bodyweight daily for 5 days. Do not mix with other drinking water medications (Note that squabs on crop milk cannot be effectively medicated by drinking water medication of parent birds).

ADMINISTRATION

Birds must have access only to medicated water. Ensure drinking system is kept clean before, during and after medication. Medication should be continuous for 3 to 5 days (except for chickens and turkeys weigh more than 1kg) according to severity of disease.

A fresh solution should be made each day. Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withdrawal period.

9. Mode of Administration:

Oral

10. Contraindications:

DO NOT USE in birds which are producing or may in the future produce eggs or egg products which may be used or processed for human consumption.

11. Warning & Precautions:

Special warnings for each target species

None, according to available data

Special precautions for use

i) Special precautions for use in animals

None.

ii) Special precautions to be taken by the person administering the product to animals

Persons mixing or handling this product should avoid inhalation of any dust and contact with skin. Wear either a disposable half-mask respiratory conforming to European Standard EN149 or a non-disposable respirator to EN140 with filter EN143.

Rubber gloves should be worn when mixing or handling this product.

Contaminated clothing should be removed and any splashes to the skin or eyes should be washed immediately.

Hands and exposed skin should be washed thoroughly after use.

Sulphonamides may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to sulphonamides may lead to cross reaction with other antibiotics. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitive to sulphonamides.
2. If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning.

12. Drug Interactions:

None known

13. Pregnancy & Lactation:

DO NOT USE in birds which are producing or may in the future produce eggs or egg products which may be used or processed for human consumption.

14. Side effects:

None recorded.

15. Symptoms & Treatment of Overdose:

No treatment specified.

16. Withdrawal Period:

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

EGGS: DO NOT USE in birds which are producing or may in the future produce eggs or egg products which may be used or processed for human consumption.

17. Storage Conditions:

Store below 30° C in dry place, protected from light.

18. Dosage Form:

Powder in the form of powder to be dissolved in water.

19. Packaging:

- 1) Pack of 1 kg in HDPE Plastic Container
- 2) Pack of 500 g in HDPE Plastic Container
- 3) Pack of 1 kg pack in Aluminium Sheet Lacquered
- 4) Pack of 500 g Pack in Aluminium Sheet Lacquered

20. Disposal of Container:

To be disposed as per local authority requirement.

21. Expiry date:

2 years from date of manufacturing.

22. Shelf-life after reconstitution or dilution:

To be used within 24 hours after dilution.

Manufactured & Distributed by:

Pelangi Vet Pharma Sdn Bhd. (665097-K)

Lot 8, PERDA Industrial Park,
Lorong IKS, Simpang Ampat B,
14100 Simpang Ampat, S.P.S,
Pulau Pinang, Malaysia.

Last update on: 13 June 2017