

BENMOXYL 500MG/G PREMIX POWDER

Composition

Each g contains:

Amoxicillin Trihydrate500mg
(equivalent to Amoxicillin 436mg)

Product Description

Benmoxyl 500mg/g Premix Powder is an off white powder

Pharmacodynamics

Amoxicillin is a semi-synthetic penicillin, derived from 6 APA (6-penicillanic acid). It is a slightly acid compound, slightly fat-soluble, stable in acid medium and can be administered by enteral or parenteral route.

Amoxicillin is a broad spectrum antibiotic with a bactericide effect primarily towards sensitive microorganisms. It causes deteriorations to the bacterial cell wall, leading to osmotic lysis of the cell.

Pharmacokinetics

After oral administration in feed, amoxicillin bioavailability is approximately 30%. The fraction reaching the blood stream is metabolized to a slight extent. The metabolite identified in laboratory species is penicilloic acid that is rapidly eliminated in the urine. Amoxicillin is weakly bound to plasma proteins (about 20%) and is widely distributed in different organs.

Indication

In pigs: Treatment of respiratory diseases due to *Streptococcus suis*, limited to reducing mortality.

Mode of Administration

Orally. Mixed into feed.

Recommended Dosage

Orally. Mixed into feed.

20 mg of amoxicillin (0.046 g BENMOXYL 500MG/G PREMIX POWDER) per kg body weight by day for 5 consecutive days, by oral route in feed.

Contraindications

Do not use in animals with serious kidney (accompanied by anuria and oliguria) and/or liver failure.

Do not used in animals with known hypersensitivity to betalactamins.

The use of the product is contraindicated when a resistance to amoxicillin is known.

Do not use in the presence of β -lactamase producing bacteria.

Do not administer to rabbits, guinea-pigs, hamsters or gerbils.

Warnings and Precautions

Special warnings

Animals with reduced feed intake and/or disturbed general condition have to be treated parenterally.

Special precautions for use

i) Special precautions for use in animals

Animals with distinctive clinical signs of streptococcosis require individual treatment.

Use of the veterinary medicinal product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to amoxicillin.

This drug premix is intended for the preparation of solid drug feed and cannot be used as is; the concentration of incorporation of the drug premix in solid feed must not be lower than 5 kg/ton.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

When feed is being prepared, avoid skin contact.

Whilst handling the product, wear a mask, coveralls, protective goggles and gloves at all times.

Avoid inhalation of dust. Wear either a disposable half-mask respirator.

To rule out any risk of ingestion it is recommended not to eat, or drink while using the product and to wash the hands after use.

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporins and vice versa. Allergic reaction to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.
2. Handle this product with great care to avoid exposure taking all recommended precautions.
3. If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments

Do not administer together with bacteriostatic anti-infectious agents (tetracyclines, sulphonamids...).

Do not use simultaneously with neomycin as it prevents the absorption of oral penicillins.

Pregnancy and Lactation

Laboratory studies have not produced any evidence of teratogenic, embryotoxic or maternotoxic effects of amoxicillin. Nevertheless, no study was performed in the target species during pregnancy or lactation. Therefore, use only accordingly to the benefit/risk assessment by the responsible veterinarian.

Adverse Effects/Undesirable Effects

Gastrointestinal signs may sometimes be observed (diarrhea).

Penicillins may cause allergic reactions after the administration of the product. Allergic reactions to these substances may sometimes be severe (anaphylaxis).

Overdose and Treatment

A part from allergic reactions, penicillins have minimal direct toxicity.

There is no antidote, in case of accidental overdosage, the treatment is symptomatic.

Withdrawal Period(s)

Meat and offal: 5 days

Storage condition

Store in a cool dry place, protected from sunlight. Do not store above 30°C.

Keep out of reach of children/Jauhkan dari kanak-kanak.

Shelf life

Shelf life as packaged for sale: 12 months

Shelf life after reconstitution into feed: 24 hours

Pack size(s)

Paper bag of 10kg and 20kg

Product Registration Holder/ Manufacturer

Osmosis Nutrition Sdn. Bhd. (594876A)

Lot 16014(P.T. 24341), Jalan Nilam 3,

Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan, Malaysia.

Tel: +606-799 9090 Fax:+606-799 9155

Date of revision of Package Insert

13/03/2024