

OXYCEN 200 mg/ml Solution for injection

COMPOSITION PER ML

Oxytetracycline (as dihydrate) 200 mg
Excipients, q.s.

PRODUCT DESCRIPTION

Clear and slightly yellow liquid.

PHARMACODYNAMICS / PHARMACOKINETICS / ENVIRONMENTAL PROPERTIES

Pharmacotherapeutic group: Antibacterial medication for systemic use. Tetracyclines.
ATCvet code: QJ01AA06

Pharmacodynamic properties: Oxytetracycline is a bacteriostatic antibiotic that inhibits protein synthesis in susceptible bacteria. Inside the cell it binds irreversibly to receptors on the 30s subunit of the bacterial ribosome where it interferes with the binding of the amino-actyl transfer RNA to the acceptor site on the messenger RNA ribosome complex. This effectively prevents the addition of the amino acids to the elongating peptide chain, inhibiting protein synthesis.

Pharmacokinetic properties: Following intramuscular injection, peak blood levels are achieved within 4-8 hours and persist for at least four days.

TARGET SPECIES

Cattle, sheep and pigs

INDICATIONS

The product is indicated for the treatment of infections caused by oxytetracycline susceptible bacteria in cattle, sheep and pigs as follows:

Cattle: - Pasteurellosis and respiratory tract infections caused by *Mannheimia haemolytica* or *Pasteurella multocida*.

- Umbilical infections and septic arthritis caused by *Trueperella pyogenes*, *Escherichia coli* or *Staphylococcus aureus*.

- Clinical Mastitis caused by *Trueperella pyogenes*, *Escherichia coli*, *Staphylococcus aureus*, *Streptococcus agalactiae* or *Streptococcus uberis*.

- Metritis caused by *Escherichia coli*

Sheep: - Pasteurellosis and respiratory tract infections caused by *Mannheimia haemolytica* or *Pasteurella multocida*.

- Umbilical infections and septic arthritis caused by *Trueperella pyogenes* or *Escherichia coli*.

- Clinical Mastitis caused by *Trueperella pyogenes*, *Escherichia coli* or *Staphylococcus aureus*.

- Erysipelas caused by *Erysipelothrix rhusiopathiae*.

- The product can also be used for treatment and metaphylaxis of enzootic abortion in sheep caused by *Chlamydophila abortus*.

Pigs: - Pasteurellosis and respiratory tract infections caused by *Mannheimia haemolytica* or *Pasteurella multocida*.

- Umbilical infections and septic arthritis caused by *Trueperella pyogenes*, *Escherichia coli* or *Staphylococcus aureus*.

- Clinical Mastitis caused by *Escherichia coli*.

- Erysipelas caused by *Erysipelothrix rhusiopathiae*.

- Atrophic rhinitis caused by *Bordetella bronchiseptica* or *Pasteurella multocida*.

DOSAGE AND METHOD OF ADMINISTRATION

The product is to be administered by deep intramuscular injection. The recommended dose rate is 20 mg/kg bodyweight (i.e. 1 ml per 10 kg bodyweight). The product is recommended for a single administration only. The cap may be safely punctured up to 30 times. When treating groups of animals, use a draw-off needle.

Maximum volume to be administered per injection site:

Cattle: 20ml, Pigs: 10ml, Sheep: 5ml.

CONTRAINDICATIONS

Not recommended for use in horses, dogs or cats. Do not use in animals with hepatic or renal damage. Do not use in known cases of hypersensitivity to the active substance or to any of the excipients.

SPECIALS WARNING FOR EACH TARGET SPECIES

None.

SPECIAL PRECAUTIONS FOR USE

i) Special precautions for use in animals: Do not dilute. If concurrent treatment is administered use a separate injection site.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals: Wash hands after use. In case of contact with eyes or skin, wash immediately with water as irritation may occur.

INTERACTIONS WITH OTHER MEDICINES

It is not recommended to administer bacteriostatic and bactericidal antimicrobials concurrently. The tetracyclines are known for their affinity and tendency to form insoluble complexes with divalent and trivalent cations, especially calcium, iron, magnesium and copper.

USE DURING PREGNANCY AND LACTATION

The product can be safely administered to lactating animals. The active substance, oxytetracycline, readily crosses the placenta and concentrations in the foetal blood may reach those of the maternal circulation, although the concentration is usually somewhat lower. Tetracyclines are deposited in teeth, causing discolouration, enamel hypoplasia and reduced mineralisation. Tetracyclines can also retard foetal skeletal development. As such, the product should only be used in the last half of pregnancy following risk benefit assessment by the responsible veterinarian. Oxytetracycline is excreted in milk; concentrations are generally low.

ADVERSE EFFECTS/UNDESIRABLE EFFECTS

Although well tolerated occasionally a slight local reaction of a transient nature may be observed. Tetracyclines have also been associated with photosensitivity reactions and, rarely, hepatotoxicity and blood dyscrasias. Oxytetracycline given to young animals can cause a yellow, brown or grey discolouration of bones and teeth. High dose or chronic administration may delay bone growth or healing.

SYMPTOMS AND TREATMENT OF OVERDOSE

There is no known specific antidote, if signs of possible overdose occur, treat the animal symptomatically.

INCOMPATIBILITIES

Refer to section Interactions with other medicines.

WITHDRAWAL PERIODS

Cattle: Meat: 31 days. Milk: 10 days.

Sheep: Meat: 9 days. Milk: 7 days.

Pigs: Meat: 18 days.

SHELF-LIFE

Shelf-life: 2 years from the manufacturing date. Shelf-life after first opening of the container: 28 days.

STORAGE CONDITIONS

Store below 25°C and protect from moisture.

PACK SIZES

Vials containing 100 ml and 250 ml. Pack with 10 vials x 100 ml and 10 x 250 ml.

DATE OF REVISION OF PACKAGE INSERT

03/06/2020

JAUHI DARI KANAK-KANAK**NAME & ADDRESS OF REGISTRATION HOLDER**

Yenher Agro-Products Sdn. Bhd. / 1628, Jalan IKS Simpang Ampat 1, Taman IKS Simpang Ampat, 14100 Simpang Ampat, Seberang Perai Selatan, Pulau Pinang, Malaysia.

NAME & ADDRESS OF MANUFACTURER

S.P. VETERINARIA, S.A.

Ctra. Reus-Vinyols, Km. 4,1

43330 Riudoms/Spain