



TEFIN Terbinafine HCl Spray 10mg/g

NAME AND STRENGTH OF ACTIVE SUBSTANCE:

Terbinafine Hydrochloride 1% w/w

PRODUCT DESCRIPTION:

A clear colourless to straw colour liquid with characteristic odour.

ROUTE OF ADMINISTRATION:

Topical

PHARMACODYNAMICS:

Terbinafine Hydrochloride is an allylamine which has a broad spectrum of antifungal activity in skin fungal infections such as tinea pedis (athlete's foot), tinea cruris (jock itch) and ringworm caused by dermatophytes such as *Trichophyton* (e.g. *T. rubrum*, *T. mentagrophytes*, *T. verrucosum*, *T. violaceum*), *Microsporum canis* and *Epidermophyton floccosum*. At low concentrations, Terbinafine Hydrochloride is fungicidal against dermatophytes and moulds. The activity against yeasts is fungicidal (e.g. *Pityrosporum orbiculare* or *Malassezia furfur*) or fungistatic, depending on species. Terbinafine Hydrochloride interferes specifically with fungal sterol biosynthesis at an early step. Terbinafine Hydrochloride inhibits ergosterol synthesis by inhibiting the fungal squalene monooxygenase (squalene 2,3-epoxidase), an enzyme that is part of the fungal cell wall synthesis pathway. Terbinafine Hydrochloride does not influence the metabolism of hormones or other drugs.

PHARMACOKINETICS:

Less than 5% of the dose is absorbed after topical application to humans; systemic exposure is thus very slight. At the level of the stratum corneum, fungicidal concentrations of Terbinafine Hydrochloride can be measured up to 7 days after the last administration.

INDICATIONS:

For the topical treatment of fungal infections

CONTRAINDICATIONS:

Hypersensitivity to Terbinafine Hydrochloride, or any component in the preparation.

ADVERSE EFFECTS:

Local symptoms such as pruritus, skin exfoliation, application site pain, application site irritation, pigmentation disorder, skin burning sensation, erythema and scab may occur at the site of application. These minor symptoms must be distinguished from hypersensitivity reactions such as widespread pruritus, rash, bullous eruptions and hives, which are reported in sporadic cases and require discontinuation. In case of accidental contact with the eyes terbinafine hydrochloride may be irritating to the eyes. In rare cases the underlying fungal infection may be aggravated. Adverse reactions are listed below by system organ class and frequency. Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100$ to $<1/10$); uncommon ($\geq 1/1,000$ to $<1/100$); rare ($\geq 1/10,000$ to $<1/1,000$); very rare ($<1/10,000$), or not known (cannot be estimated from available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. **Immune system disorders.** Not known: Hypersensitivity. **Eye disorders.** Rare: Eye irritation. **Skin and subcutaneous tissue disorders.** Common: Skin exfoliation, pruritus. Uncommon: Skin lesion, scab, skin disorder, pigmentation disorder, erythema, skin burning sensation. Rare: Dry skin, dermatitis contact, eczema. Not known: Rash. **General disorders and administration site conditions,**

Uncommon: Pain, application site pain, application site irritation.

Rare: Condition aggravated.

DRUG INTERACTION:

Unknown.

PRECAUTIONS/WARNINGS:

TEFIN Terbinafine HCl Spray 10mg/g should be used with caution in patients with lesions where alcohol could be irritating. TEFIN Terbinafine HCl Spray 10mg/g is for external use only. It may be irritating to the eyes. TEFIN Terbinafine HCl Spray 10mg/g should not be used on the face. In case of accidental contact with the eyes, rinse eyes thoroughly with running water. Avoid inhalation. In case of accidental inhalation, consult a physician if any symptoms develop or persist. TEFIN Terbinafine HCl Spray 10mg/g contains propylene glycol, which may cause skin irritation.

PREGNANCY AND LACTATION:

Usage of TEFIN Terbinafine HCl Spray 10mg/g by pregnant women and those who are trying to become pregnant, consult doctor or pharmacist for further advice. Do not use the spray during breast-feeding period.

RECOMMENDED DOSAGE:

Apply TEFIN Terbinafine HCl Spray 10mg/g to the affected areas once a day, for one week. Spray from a distance of 5cm to 10cm. Regular applications is essential for successful treatment. Treatment should be continued for two weeks after all clinical signs have disappeared. Attention to hygiene is important in the management of fungal diseases, particularly of the folds and the feet. If no improvement is seen after 4 weeks, diagnosis should be reviewed.

OVERDOSAGE:

The low systemic absorption of topical terbinafine renders overdosage extremely unlikely. When a large amount of TEFIN Terbinafine HCl Spray 10mg/g be inadvertently ingested, the adverse effects to be expected include headache, nausea, epigastric pain and dizziness. In case of accidental oral ingestion, the alcohol content has to be considered. TEFIN Terbinafine HCl Spray 10mg/g contains 25% w/w alcohol.

TREATMENT OF OVERDOSE:

If accidentally ingested, the recommended treatment of overdosage consists of eliminating the active substance, primarily by the administration of activated charcoal, and giving symptomatic supportive therapy if needed.

PACKING:

15mL in HDPE bottle.

STORAGE:

Store below 30°C. Keep out of reach of children.

Jauhi daripada kanak-kanak.

Recommended shelf-life: 2 years.

MANUFACTURED BY:

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,
42000 Port Klang, Selangor, MALAYSIA.

Date of revision: JUNE 2023