

Front

TRIMIC™
Metronidazole & Miconazole
Nitrate Vaginal Suppositories

Composition:
Each vaginal suppository contains:
Metronidazole BP 500 mg
Miconazole Nitrate BP 100 mg
Excipients q.s.

Description: White to off white Opaque bullet shaped suppositories.

Pharmacodynamics
Trimic (Metronidazole, Miconazole Nitrate vaginal suppository) contains miconazole for antifungal and metronidazole for antibacterial and antitrichomonal effects. Miconazole nitrate which is an antifungal has a wide spectrum of activity and is particularly effective against pathogen fungi including *Candida albicans*. In addition, miconazole nitrate is effective against Gram (+) bacteria. Metronidazole is an antibacterial and antiprotozoal agent that is effective against *Gardenella vaginalis* and against anaerobic bacterial including anaerobic streptococci and *Trichomonas vaginalis*.

Pharmacokinetics
Absorption of miconazole nitrate through the intravaginal route is very low. Bioavailability of metronidazole by this route is 20 percent compared to the oral route. Miconazole nitrate cannot be detected in plasma after vaginal application of Trimic. Steady state level of metronidazole in plasma reaches 1.6–7.2 µg/mL. Metronidazole is metabolized in the liver. The hydroxyl-metabolite is effective. The half-life of metronidazole is 6-11 hours. Approximately 20 percent of the dose is excreted unchanged in urine.

Indications
TRIMIC (Metronidazole & Miconazole Nitrate Vaginal Suppositories) should be used in the treatment of vaginal candidiasis, bacterial and trichomonal vaginitis, or vaginitis due to mixed infections.

Dosage and administration:
Do not use without consulting a physician.
If it is not advised otherwise by a physician, 1 pessary should be inserted high into the vagina at night for 14 days or 1 pessary at night and 1 pessary in the morning for 7 days. In recurrent cases, or when the vaginitis has been resistant to other

treatments, 1 pessary should be inserted high into the vagina at night and in the morning for 14 days.
For elderly patients (over 65 years): As for younger adults.
Children: Not recommended for children.
Not to be swallowed or not to be applied by other routes.

Contraindications
You should not use Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories) if:
- You suffer from hypersensitivity to any of the ingredients of the suppository
- You are pregnant (particularly in the first 3 months)
Your doctor needs to carefully evaluate whether you may use Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories) if you suffer from:
- Severe problems with your liver (liver function disorders including metabolic disease called "porphyria")
- Diseases of the nervous system
- Disturbances in the production of blood (hematopoiesis)

WARNINGS/PRECAUTIONS
You must not take alcohol during the therapy and for at least 24-48 hours after the end of course of treatment because of the possibility of increased alcohol incompatibility. You must not use the suppository together with the contraceptive diaphragms and condoms since the vaginal suppository base may affect the rubber, which may possibly break.

Sila dapatkan nasihat doktor atau ahli farmasi sebelum menggunakan keluaran ini jika anda mengambil ubat warfarin, iaitu sejenis ubat antipembekuan darah, kerana lebam/ pendarahan pada gusi/ hidung boleh berlaku secara spontan.

(Please consult your physician/ pharmacist before using this product if you are on the anticoagulant medicine warfarin, because bleeding from nose/ gums or bruising may occur spontaneously).

Pregnancy and Lactation
After the first three months of pregnancy, Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories) can be used when considered essential by your doctor, but must be used under observation. You must not breastfeed your baby while using Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories), since active ingredients pass into the milk. You may start breast feeding again after 24-48 hours after the treatment is finished.
Interactions with other medications
Due to absorption of active ingredients of Trimic (Metronidazole & Miconazole Nitrate Vaginal

Back

Suppositories) interactions may occur if you used the preparation with alcohol. Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories) may lead to an increased alcohol incompatibility) or with any of the following drugs (inform your doctor if you take them):
- Drugs that make your blood thinner (oral anti coagulants)
- Anti-epileptics (phenytoin, phenobarbital)
- A drug reducing the acid in your stomach (cimetidine)
- A drug for the treatment of alcoholism (disulfiram)
- A drug against psychosis and depression (lithium)
- Anti-allergic medicines (astemizole and terfenadine)
Laboratory tests may change while you use Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories):
Interference with blood levels of liver enzymes, glucose (hexokinase method), theophylline, and procainamide have been observed. If blood tests are made, remind your doctor that you use Trimic.

Adverse recations
Medical events judged to be common, uncommon and rare are listed below.

Vaginal disorders
Vaginal irritation (burning, itching) owing to inflammation of vaginal mucosa in vaginitis, vaginal irritation can be seen after the administration of the first ovule or towards the third day of treatment. These complaints disappear quickly when the treatment continues. If there is severe irritation, contact your doctor because it may be necessary to stop the treatment.

Stomach and intestinal disorders
Abdominal or cramp, Metallic taste, constipation, dry mouth, seldom diarrhea, lack of appetite, vomiting and nausea.
Nervous system disorders
Headache, movement disorders (ataxia), dizziness, mental changes, disorders of the nerves (peripheral

neuropathy) after overdose or long period of usage, cramps.

Skin disorders
Rash

Blood disorders
Decrease of white blood cells

Overdosage
There is no human experience with overdose of Trimic (Metronidazole & Miconazole Nitrate Vaginal Suppositories). If large quantities of the product are accidentally taken, an appropriate method of emptying your stomach may be considered necessary. Symptoms of Metronidazole overdose are nausea, vomiting, abdominal pain, diarrhea, itching, metallic taste, movement disorders (ataxia), vertigo, unusual sensations on the skin (paresthesia), cramps, decrease of white blood cells, and darkening of urine. Symptoms of Miconazole nitrate overdose are nausea, vomiting, sore throat and mouth, decrease appetite, headache and diarrhea.

Storage condition
Store in dry place, below 30°C.
Protect from light.
Keep the product in its original package.
Keep out of reach and sight of children.

Dosage forms & Packing available:
Two blister strips of 7 vaginal suppositories are packed in a carton along with an insert.

Date of revision of PI: June 2018.



13005146

Manufactured by :
BLISS GVS PHARMA LTD.
Plot No. 11, Survey No.38/1, Dewan Udyog Nagar, Aliyali Village, Taluka Palghar, Thane 401 404, Maharashtra State, India.

Product Registration Holder:
Healol Pharmaceuticals Sdn. Bhd., 74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju, 53300 Kuala Lumpur, Malaysia.