

Veterinary Package Insert

TP-Amox 500 WSP Amoxicillin Trihydrate 500 mg/g Powder

PRODUCT DESCRIPTION

A white, off-white to light yellow colour powder, which contains 500 mg/g of amoxicillin trihydrate in 100g, 500g or 1kg sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against Gram positive and Gram negative bacteria. It owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

INDICATION

Treatment of infections in chickens and ducks caused by bacteria susceptible to amoxicillin. Not effective against beta-lactamase producing organisms.

Pigs: For the treatment of salmonellosis and pasteurellosis.

TARGET SPECIES

Chickens, ducks, pigs

RECOMMENDED DOSAGE AND ADMINISTRATION

Chicken and ducks:

The product is administered in the drinking water. Prepare the solution with fresh tap water immediately before use. Once opened, use the contents of one sachet immediately. Any unused medicated water which should be discarded after 12 hours.

In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

The following formula may be used to calculate the amount of product required per day (in grams):

Number of birds x average live weight (kg)
25 (for 20 mg/kg) or 33 (for 15 mg/kg)

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid under-

dosing. The uptake of medicated water depends on the clinical condition of the birds. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake.

Chickens:

The recommended dosage is 15 mg amoxicillin trihydrate per kg bodyweight. The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks:

Recommended dosage is 20 mg amoxicillin trihydrate per kg bodyweight for 3 consecutive days.

Pigs:

For the medication of Pigs, TP-Amox 500 WSP Amoxicillin Trihydrate 500mg/g Powder may be administered in the drinking water or administered by addition to liquid feeds produced with commercial feed. It may not be used in dry feeds.

i) For administration via the drinking water, dose of amoxicillin trihydrate 20mg/kg body weight daily, should be divided and administered at approximately 12 hourly intervals for up to 5 days. Dissolve the content of one sachet (100g of TP-Amox 500 WSP Amoxicillin Trihydrate 500mg/g Powder) in appropriate volume of water required immediately before use. Any medicated water not consumed within 12 hours should be discarded.

ii) For administration via liquid feed, dose of amoxicillin trihydrate 20mg/kg body weight daily, should be given up to 5 days. Medicated liquid feed should be freshly prepared on at least 3 occasions per day over the 5-day treatment period. Dissolve the required amount of TP-Amox 500 WSP Amoxicillin Trihydrate 500mg/g Powder to give a fluid at 25g/L of the product before adding to dry feed. This may be done by dissolving the content of one sachet (100g of TP-Amox 500 WSP Amoxicillin Trihydrate 500mg/g Powder) in approximately 4L of water immediately before use.

The medicated liquid feed should not be fermented and should not be stored. Stability of amoxicillin in all commercial feeds has not been established. In order to ensure that any loss of amoxicillin activity is minimized, the quantity of medicated liquid feed prepared should not exceed the amount of feed which will be consumed within 4 hours.

CONTRAINDICATIONS

The product should not be administered to rabbits, hamsters, gerbils and guinea pigs, or to birds producing eggs for human consumption.

WARNINGS & PRECAUTIONS.

Special warnings for each target species: None.

(i) Special precautions for use in animals

Pigs: The uptake of medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed or water, animals should be treated parenterally. Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1) Do not handle this product if you know you are sensitized or if you have been advised not to work with such preparations.

2) Handle this product with great care to avoid exposure, taking all recommended precautions.

3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

- Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143

- Wear gloves during preparation and administration of medicated water or liquid feed

- Wash any exposed skin after handling the product or medicated water or feed

- Wash hands after use

INTERACTIONS WITH OTHER MEDICATIONS

None known for drugs use in animals.

PREGNANCY AND LACTATION

Use only according to the benefit/risk assessment of the responsible veterinarian.

ADVERSE EFFECTS

Not known.

OVERDOSE & TREATMENT

No problems with overdose have been reported. Treatment should be symptomatic and no specific antidote is available.

WITHDRAWAL PERIOD

Chickens (meat & offal): 1 day

Ducks (meat & offal): 9 days

Pigs (meat & offal): 2 days

Note: Amoxicillin is not permitted for use in birds producing eggs for human consumption.

MAXIMUM RESIDUAL LIMIT (MRL)

All food producing species

Muscle: 50 µg/kg

Liver: 50 µg/kg

Kidney: 50 µg/kg

Fat: 50 µg/kg

STORAGE CONDITION

Store in a cool, dry place (below 30°C). Protect from sunlight.

SHELF LIFE

3 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

THYE PHARMA SDN. BHD.

No.2 Lorong Industri Ringan Permatang
Tinggi 8,
Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised November 2018