

NEOTM
SOLUBLE POWDER
(FOR VETERINARY USE ONLY)

COMPOSITION

Neomycin (as sulfate) 200 mg/g
Oxytetracycline 200 mg/g

PRODUCT DESCRIPTION

The finished product is light yellowish color powder.

TARGET SPECIES

Cattle (calves), sheep (lambs), swine and poultry.

PHARMACODYNAMIC

Neomycin is an antibiotic of the aminoglycoside obtained from Streptomyces fradiae. Its spectrum of activity includes Gram positive, especially staphylococci and less active streptococci, and Gram negative, especially Escherichia coli. Neomycin binds to the 30S subunit of the bacterial ribosome, which disturbs the reading of the constituent code of the messenger RNA, and finally the bacterial protein synthesis. At high concentrations, it has been shown that aminoglycosides damage the bacterial cell wall, adding bactericidal properties to bacteriostatic properties.

Oxytetracycline binds reversibly to receptors of the 30S ribosomal fraction, this leads to a blocking of the binding of aminoacyl-tRNA to the corresponding site of the ribosome complex messenger-RNA. This results in inhibition of protein synthesis and therefore a growth arrest of the bacterial culture. Oxytetracycline is a bacteriostatic. The bacteriostatic activity of oxytetracycline involves penetration of the substance into the bacterial cell. The penetration of oxytetracycline is exercised by both passive and active dissemination. The main possible resistance mode is related to the presence of an R factor responsible for a decrease in active transportation oxytetracycline. Oxytetracycline is a broad spectrum antibiotic. It is primarily active against microorganisms Gram positive and negative, aerobic and anaerobic, and against mycoplasma, Chlamydia and Rickettsiae. The acquired resistance to oxytetracycline was reported. Such resistance is usually plasmid origin. Crossresistance to other tetracyclines is possible. Continuous treatment with low doses of oxytetracycline can also lead to increased resistance to other antibiotics.

PHARMACOKINETIC

Neomycin is very poorly absorbed from the gastrointestinal tract. In plasma and tissue concentrations of neomycin are very low. It is present in large amount, and persistently in the different sections of the digestive tract. Neomycin is almost exclusively eliminated in the faeces after oral administration in calves, the absorption is very low (111%) with 90% of the dose excreted in the faeces of which 70 to 80% present as unchanged.

For the most species, oxytetracycline is quickly (2 to 4 hours) absorbed after oral administration in animals fasted and its bioavailability is between 60% and 80%. The bioavailability may decrease in the presence of food in the stomach because oxytetracycline forms insoluble chelates with divalent or trivalent cations (Mg, Fe, Al, Ca) they contain.

The oxytetracycline is bound to plasma proteins variably depending on the species (20-40%). Its distribution is wide. Oxytetracycline diffuses throughout the body, the highest concentrations found in the kidneys, liver, spleen and

lungs. Oxytetracycline crosses the placenta.

Oxytetracycline is excreted unchanged mainly via urine. It is also excreted in the bile, but a large percentage of oxytetracycline is reabsorbed by the small intestine (enterohepatic cycle).

INDICATIONS

For calf, lamb, pigs and poultry:

Treatment of gastrointestinal infections germs sensitive to oxytetracycline and neomycin.

WARNINGS AND PRECAUTIONS

Special warnings for each target species:

This powder for oral solution is intended to target species described under "Target species".

Special precautions for use:

i. Special precautions for use in animals:

This powder for oral solution is intended to be dissolved in milk, the liquid food or drinking water and cannot be used as it is.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not handle this product in case of known allergy to tetracycline or aminoglycoside.

Hypersensitivity to neomycin and other aminoglycosides can be crossed.

Allergic reactions to these substances may be severe.

In case of hypersensitivity, avoid contact with the product.

Handle it by taking the recommended precautions to avoid any risk of exposure.

Wearing a mask, goggles and protective gloves is recommended when preparing the solution. Avoid breathing dust and avoid contact with skin and eyes.

In case of appearance of skin rash, take medical advice.

In case of occurrence of edema of the face, lips, eyes or if they develop difficulty breathing, seek medical attention immediately.

In case of accidental projection into the eyes, rinse thoroughly with clean water.

Wash hands after use of the product.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

CONTRAINDICATIONS

Do not use in case of:

- Known hypersensitivity to tetracyclines and / or aminoglycosides,
- Renal insufficiency,
- Known resistance to tetracyclines and / or aminoglycosides.

ADMINISTRATION & DOSAGE

For oral administration

Calves, lambs, pigs:

10 mg of neomycin and oxytetracycline 10 mg per kg body weight every 12 hours orally for 3 days, 5 g of powder per 100 kg body weight morning and evening in milk, milk replacer, the liquid food or drinking water for 3 days.

Poultry:

20 mg of neomycin and oxytetracycline 20 mg per kg body weight per day orally for 3 days, 10 g of powder per 100 kg bodyweight per day for 3 days in drinking water.

INTERACTION WITH OTHER MEDICAMENTS

Divalent or trivalent cations (Mg, Fe, Al, Ca) can chelate tetracycline.

Tetracycline should not be administered with antacids, gels based on aluminum, preparations containing vitamins or minerals as insoluble complexes are formed, which reduces the absorption of the antibiotic.

PREGNANCY AND LACTATION

The oxytetracycline showed no sign of embryotoxicity or teratogenicity in laboratory animals.

In mammals, oxytetracycline through the placenta, causing tooth staining and slower fetal growth.

The tetracyclines are found in breast milk. The safety of the product has not been evaluated in pregnant and lactating animals. Use of the product in pregnant and lactating animals will be an evaluation of the benefit / risk balance by the veterinarian.

OVERDOSE AND TREATMENT

Refer to "SIDE EFFECT"

SIDE EFFECT

Like all tetracycline, oxytetracycline can trigger side effects such as gastrointestinal disorders and less frequently allergic reactions and photosensitivity.

WITHDRAWAL TIME

Cattle and sheep:

Meat and offal: 14 days.

Milk: See "Target Species" (the specialty is not intended for dairy milk producing females).

Pigs:

Meat and offal: 14 days.

Poultry:

Meat and offal: 14 days.

Eggs: zero days.

SHELF LIFE: 36 months

SHELF LIFE after first opening & reconstitution: 24 hours

STORAGE

Store in temperature not exceeding 30°C. Strictly avoid light & heat exposure.

PACKAGING

The powder filled 500 g in aluminum foil.

DATE OF REVISION: 31 May 2018



Product registration holder & Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,

Taman Perindustrian USJ 1,

47610 Subang Jaya, Selangor.