

PDP-ALBENDAZOLE SUSPENSION

Composition:

Each liter contains.

Albendazole BP Vet 19.00 gm

Preservative:

Sodium Methyl Paraben BP 0.2 %

Sodium Propyl Paraben BP 0.02 %

Sodium Benzoate BP 0.2 %

Description: A light orange coloured suspension

Pharmacology:

Albendazole's mode of action has been shown to be based on the inhibition of the uptake and efficiency of utilization of glucose by roundworms and tapeworms and similar activity is presumed in liver fluke

Albendazole mode of action is to inhibit energy metabolism by uncoupling phosphorylation reactions. In fasciola hepatica, cellular changes due to Albendazole include deformed mitochondria, dilatation of the rough endoplasmic reticulum, vesiculation of the golgi field and a reduction or absence of secretory granules.

Pharmacokinetics:

The pharmacokinetics of albendazole are the same as when the products are administered separately. Peak plasma levels are attained 15 hours after treatment. The depletion curve is linear and indicates a drug half-life 10 hours.

Albendazole absorbed from the gut and rapidly metabolized in the liver almost completely to the 2- amino sulfone metabolite plus 3 or 4 other metabolites. Peak plasma albendazole levels are reached within 8 to 24 hours. Due to plasma binding, levels persist which are adequate to exert a sustained anthelmintic effect on Haemonchus contortus. Tissue residues, however, are minimal.

Albendazole is metabolized primarily in the liver and, due to concentration in the bile, excretion of metabolites is largely via the faecal route.

Indications:

Control of susceptible mature and immature gastrointestinal roundworms, large lung worms, liver fluke, nasal bots; reduces output of viable worm and fluke eggs and provides sustained control of barber's pole worm in sheep.

Control of the following albendazole sensitive stomach, intestinal and bowel worms: barber's pole worm (Haemonchus contortus), stomach hair worm (Trichostrongylus axei), black scour worm (Trichostrongylus sp.), small brown stomach worm (Ostertagia sp.), small intestinal worm (Cooperia sp.), thin necked intestinal worm (Nematodirus sp) hookworm (Bunostomum trigonocephalum), nodule worm (Oesophagostomum columbianum), large bowel worm (O. venulosum), large mouthed bowel worm (Chabertia ovina), large lung worm (Dictyocaulus viviparus) and tapeworms (head and sections) (Moniezia sp.).

Albendazole is also an effective aid in the control of adult liver fluke and in reducing the output of viable worm and fluke eggs.

Dosage and Administration:

Roundworm, lungworm, tapeworm. 1 mL/5kg bodyweight orally.

Roundworm, lungworm, tapeworm, adult liver fluke. 1 mL/4kg bodyweight orally.

Control of roundworms, lungworm, tapeworm, and to reduce output of viable worm eggs: 1 ml/5 kg liveweight. Control of roundworms, lungworm, tapeworm, and to reduce output of viable worm eggs, and aid in control of adult liver fluke: 1 ml/4 kg liveweight.

Drench sheep orally using an appropriate according to the heaviest animal in the sheep to determine the heaviest animal. Gun to make sure it is delivering correct dose to the animal.

Precautions Safety directions. Harmful if absorbed by skin contact or swallowed. Avoid contact with skin and wash hands after use. First aid. If poisoning occurs, contact a doctor or Poisons Disposal. Destroy empty containers by breaking, crushing, puncturing or shredding where applicable. Do not contaminate dams, rivers or streams. Dispose of the containers at a local authority landfill that does not burn its refuse. If such a landfill is not readily available in your area, bury containers at a depth of 500 mm or more at a licensed approved disposal site. In some States wastes can only be buried at a licensed landfill. Do not burn empty containers or Product.

Do not mix with other drenches. Shake container before use.

Withholding Periods Meat Sheep, goats: 10 days. Milk. Do not use in sheep or goats which are producing or may in the future produce milk or milk products for human consumption.

Contraindications: Use within 7 days of bromsalans.

Pregnancy and lactation:

During early pregnancy, all medication should be administered with care, and it is particularly important not to exceed the recommended dosage at this time.

Side effects:

There are no known side effects of any drug with Albendazole Susp

Storage: Store in dry place, 30°C and protect from light.

Symptoms and treatment of overdose:

No report on overdose

Shelf life: 2 years from date of manufacture

Presentation: Pack of HDPE 1 Lt, 5 Lt, 25 Lt.

Manufactured by:

Pelangi Dairy Products Sdn.Bhd (665097K)
No 46, Lorong Kendi 7, Taman Merak Jaya,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia.

Date of information: Jun, 2009.