
WINAMOL Drops for Infants (Paracetamol 100mg/ml)

Name and Strength of Active Ingredient(s):

Each 1 ml contains:

Paracetamol 100 mg

Preservative:

Methyl Paraben 0.09 % w/v

Propyl Paraben 0.009 % w/v

Product Description

Red colour syrup with raspberry flavour.

Pharmacodynamics

Paracetamol has analgesic and antipyretic actions probably due to the inhibition of prostaglandin biosynthesis.

Pharmacokinetics

Paracetamol is readily absorbed from the gastro-intestinal tract and peak plasma concentrations usually occur 30 minutes to 2 hours after ingestion. Paracetamol is metabolised in the liver and largely excreted in the urine as sulphate and glucuronide conjugates. Less than 5% is excreted unchanged. The elimination half-life varies from about 1 to 4 hours.

Indications

For the relief of fever and mild to moderate pain. WINAMOL Drops is an analgesic in teething.

Recommended Dose

1-3 months (4-6 kg) : 0.6-0.9 mL (60-90mg)

3-6 months (6-8 kg) : 0.9-1.2 mL (90-120mg)

6-12 months (8-10 kg) : 1.2-1.5 mL (120-150mg)

1-2 years (10-12 kg) : 1.5-1.8 mL (150-180mg)

- For an accurate dosing, weigh your child and work out the dosage to be given based on a paracetamol 15 mg for every kg body weight.
- Do not give more than 4 doses in any 24 hour period.
- Leave at least 4 hours between doses.
- Do not give this medicine to your child for more than 3 days without speaking to your doctor or pharmacist.

Route of Administration

Oral

Contraindications

Contra-indicated in patients with a known hypersensitivity to paracetamol or any of the other constituents.

Warning and Precautions

WARNING

This preparation contains PARACETAMOL.
Do not take any other paracetamol containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.

These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Consult the physician before use if the child has liver or kidney problems and if the child's symptoms do not improve. Do not exceed the stated dose.

WINAMOL Drops for Infants contains paracetamol. It should not be used with other paracetamol-containing products.

Interactions with Other Medicaments

Before taking WINAMOL Drops for Infants, make sure to consult the physician if the patient is taking warfarin or similar medicines used to thin the blood.

Pregnancy & Lactation

Epidemiological studies in human pregnancy have shown no ill effects due to paracetamol used in the recommended dosage, but patients should follow the advice of their doctor regarding its use.

Paracetamol is excreted in breast milk but not in a clinically significant amount.

Available published data do not contra-indicate breast feeding.

Side Effects

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Stop taking WINAMOL Drops for Infants and inform the physician immediately if the child: Experience allergic reactions eg, skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face; experience a skin rash or peeling, or mouth ulcers; have previously experienced breathing problems with aspirin or nonsteroidal anti-inflammatory drugs, and experience a similar reaction with WINAMOL Drops for Infants; experience unexplained bruising or bleeding, these reactions are rare.

Symptoms and Treatment of Overdose

Early symptoms of paracetamol overdosage include pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning hepatic failure may progress to encephalopathy, coma and death. Acute renal failure with acute tubular necrosis may develop even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported. Liver damage is possible in adults who have taken 10g or more of paracetamol. It is considered that excess quantities of a toxic metabolite (usually adequately detoxified by glutathione when normal doses of paracetamol are ingested), become irreversibly bound to liver tissue.

Immediate treatment is essential in the management of paracetamol overdose. Despite a lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention. Any patient who has ingested around 7.5g or more of paracetamol in the preceding 2 hours should undergo gastric lavage. Administration of oral methionine or intravenous n-acetylcysteine which may have a beneficial effect up to at least 48 hours after the overdose, may be required. General supportive measures must be available.

Storage Condition

Keep the bottle tightly closed protected from light. Store below 30°C.

Presentation:

15ml in Amber glass bottle with white HDPE screw cap and in Amber PET bottle with white HDPE screw cap.

Registration Number:

MAL19086013XZ

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia

Date of Revision:

02 December 2024