

强康生

QIANG KANG SHENG

Amoxicillin 30%w/w Soluble Powder

(FOR VETERINARY USE ONLY)

COMPOSITION

Amoxicillin (as Amoxicillin trihydrate) 300 mg/g

PRODUCT DESCRIPTION

This product is white-like powder.

TARGET SPECIES

Chickens

PHARMACODYNAMICS

Amoxicillin is a time-dependent bactericidal antibiotic which acts by inhibiting the synthesis of bacterial cell walls during bacterial replication. It inhibits the formation of bridges between the chains of linear polymers constituting the peptidoglycan cell wall of Gram positive bacteria.

Amoxicillin is a broad-spectrum penicillin. It is also active against a limited range of Gram negative bacteria on which the outer layer of the bacterial cell wall is composed of lipopolysaccharide and proteins.

There are three main mechanisms of resistance to beta-lactams: beta-lactamase production, altered expression and/or modification of penicillin binding proteins (PBP), and decreased penetration of the outer membrane. One of the most important is the inactivation of penicillin by beta-lactamase enzymes produced by certain bacteria. These enzymes are capable of cleaving the beta-lactam ring of penicillins, making them inactive. The beta-lactamase could be encoded in chromosomal or plasmidic genes. Cross-resistance is observed between amoxicillin and other penicillins, particularly with aminopenicillins.

The use of extended spectrum beta-lactam drugs (e.g. aminopenicillins) might lead to the selection of multi-resistant bacterial phenotypes (e.g. those producing extended spectrum beta-lactamases (ESBLs)).

PHARMACOKINETICS

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids. Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

INDICATIONS

Treatment of infections in chickens caused by bacteria susceptible to amoxicillin.

CONTRA-INDICATIONS

Do not use in horses, rabbits, guinea pigs, hamsters, gerbils or any other small herbivore.

Do not use in animals with known hypersensitivity to penicillins or other β -lactam antibiotics or to any of the excipients.

ADMINISTRATION & DOSAGE

In drinking water use. Prepare the solution with fresh tap water immediately before use. Any unused medicated water should be discarded after 12 hours. In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated. The following formula may be used to calculate the required concentration of product (in milligrams of product per litre drinking water):

x mg product per kg body weight per day	X	mean body weight (kg) of animals to be treated	=	x mg product per liter drinking water
mean daily water consumption (liter) per animal per day				

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing. The uptake of medicated water depends on the clinical condition of the birds. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake. After the end of the medication period the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

The recommended dosage is 13.1 mg amoxicillin per kg bodyweight for 3 days (corresponding to 44 mg product/kg-bodyweight/day) or in severe cases for 5 days.

WARNING AND PRECAUTIONS

Special warnings for each target species:

None

Special precautions for use:

i) Special precautions for use in animals:

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Use of the product deviating from the instructions given in the PI may increase the prevalence of bacteria resistant to the amoxicillin and may decrease its effectiveness.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Avoid inhalation of dust.

Wear either a disposable half-mask respirator or a non-disposable respirator with a filter.

Penicillins and cephalosporins may cause hypersensitivity reactions which may occasionally be serious. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. In case of contact with eyes or skin, wash immediately with water. Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations. Handle this product with great care to avoid exposure, taking all recommended precautions. If you develop symptoms following exposure such as a skin rash, you should seek

medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention. Wash hands after use.

- Keep out of reach of children.
- Jauhi ubat dari kanak-kanak.

INTERACTION WITH OTHER MEDICAMENTS

The product should not be administered with antibiotics that have a bacteriostatic mode of action, such as tetracyclines, macrolides, sulphonamides.

PREGNANCY AND LACTATION

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin. Use only according to the benefit/risk assessment of the responsible veterinarian.

SIDE EFFECT

Penicillins and cephalosporins may cause hypersensitivity reactions which may occasionally be serious.

OVERDOSE AND TREATMENT

None known.

WITHDRAWAL PERIODS

Chickens (meat and offal): 1 day

The product is not authorized for use in laying birds producing eggs for human consumption and within 3 weeks of onset of laying.

SHELF LIFE: 36 months

SHELF LIFE after first opening: 24 hours

SHELF LIFE after dilution: 12 hours

STORAGE

Store at temperature not exceeding 30°C. Strictly avoid light & heat exposure.

PACKAGING

25 kg/ bag

DATE OF REVISION: 27 Jun 2019



Manufactured by:

厦门惠盈动物药业有限公司

XIAMEN HUIYING ANIMAL

PHARMACEUTICAL CO., LTD

No.3106, Guankou Road, Jimei District,
Xiamen, Fujian, China.



Product registration holder:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,
Taman Perindustrian USJ 1,
47610 Subang Jaya, Selangor, Malaysia.