

AMOSYN 500 mg/g WATER SOLUBLE POWDER

COMPOSITION:

Contains per g
Amoxicillin Trihydrate 500mg
Equivalent to Amoxicillin 436mg

DESCRIPTION

White or almost white powder; taste, slightly bitter.

PHARMACODYNAMICS:

Amoxicillin is a bactericidal antibiotic belonging to the group of semisynthetic penicillins with a broad spectrum of activity against Gram positive and Gram negative bacteria. It owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

PHARMACOKINETICS:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals

INDICATION:

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

Pigs: For the treatment of pasteurellosis

TARGET SPECIES:

Chickens, ducks, turkeys, pigs

DOSAGE:

The product is administered in the drinking water. Prepare the solution with fresh tap water immediately before use. Once opened, use the contents of one sachet immediately. Any unused medicated water should be discarded after 12 hours.

In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

The following formula may be used to calculate the amount of product required per day (in milligrams of product per litre of drinking water):

$\frac{\text{x mg product per kg bodyweight per day} \times \text{mean bodyweight (kg) of animals to be treated}}{\text{mean daily water consumption (l) per animal}} = \text{x mg product per litre drinking water}$

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing. The uptake of medicated water depends on the clinical condition of the birds. In order to obtain the correct dosage, the concentration of amoxicillin has to be adjusted taking into account water intake.

Chickens

The recommended dosage is 15 mg amoxicillin trihydrate per kg bodyweight per day (corresponding to 30 mg product/ kg body weight/ day).

The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks

Recommended dosage is 20 mg amoxicillin trihydrate per kg bodyweight per day (corresponding to 40 mg product/kg bodyweight/day) for 3 consecutive days

Turkeys

Recommended dosage is 15-20 mg amoxicillin trihydrate per kg bodyweight per day (corresponding to 30-40 mg product/kg bodyweight/day) for 3 days or in severe cases for 5 days.

Pigs

For the medication of pigs, Amosyn 500 mg/g water soluble powder may be administered in the drinking water to give 20 mg amoxicillin trihydrate per kg of bodyweight (corresponding to 40 mg product/kg bodyweight) daily for up to 5 days.

The water supply should be cleaned appropriately after the end of medication period.

CONTRAINDICATIONS:

Amosyn 500 mg/g water soluble powder should not be administered to rabbits, guinea pigs, hamsters, gerbils or any other small herbivore.

Do not use in animals with known hypersensitivity to penicillins or other β -lactam antibiotics or to any of the excipients.

Do not administer to animals with renal disease including anuria or oliguria.

WARNING AND PRECAUTION:

Special warnings for each target species

None

i. Special precautions for use in animals

Not effective against beta-lactamase producing organisms.

Pigs: The uptake of medication by animals may be altered as a consequence of illness. In case of insufficient water uptake, animals should be treated parenterally instead.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Use of the product deviating from the instructions given in the package insert may increase the prevalence of bacteria resistant to the amoxicillin and may decrease the effectiveness of the treatment.

- ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

People with known hypersensitivity to the active substance or if you have been advised not to work with such preparations should avoid contact with the veterinary medicinal product. Handle this product with great care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention. Avoid inhalation of dust.

Wear gloves during preparation and administration of medicated water or liquid feed. Wash any exposed skin after handling the product or medicated water or feed. Wash hands after use.

Container should be disposed in accordance with local requirements.

INTERACTIONS WITH OTHER MEDICATIONS

Amosyn 500 mg/g water soluble powder should not be administered with antibiotics that have a bacteriostatic mode of action, such as tetracyclines, macrolides, sulphonamides.

PREGNANCY AND LACTATION

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

Use only according to the benefit/risk assessment of the responsible veterinarian.

Do not use in birds producing eggs for human consumption or within 3 weeks of onset of laying.

SIDE EFFECTS:

Penicillins and cephalosporins may cause hypersensitivity reactions which may occasionally be serious.

SYMPTOMS AND TREATMENT OF OVERDOSE

No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

WITHDRAWAL PERIOD(S):

Meat and offal:

Chickens	1 day
Ducks	9 days
Turkey	5 days
Pigs	2 days

STORAGE

Store in a cool and dry place
Do not store above 30°C
Store in an airtight container

For Veterinary Use Only
Keep out of the reach of children
Jauhi daripada kanak-kanak

PROPOSED SHELF LIFE:

Shelf life of the veterinary medicinal product as packaged for sale: 24 months
Shelf life after first opening the immediate packaging: 3 months
Shelf life after dilution or reconstitution according to directions: 12 hours

PACK SIZE AVAILABLE:

100 g per sachet and further packed 100 sachets in one carton

Manufactured by:

Hangzhou An-I-Mate Animal Pharmaceutical Co. Ltd.

No. 2, Fengdu Road, Fengdu Industrial Garden,
Pingyao, Yuhang, Hangzhou,
China.

Marketing Authorization Holder:

Ritma Prestasi Sdn Bhd

No. 43, Jalan Meranti Jaya 11,
Taman Perindustrian Meranti Jaya,
47120 Puchong Selangor,
Malaysia

Date of Revision:

27 June 2019