

Veterinary Package Insert

TP-CTC HCL 500MG/G POWDER (50% Chlortetracycline Hydrochloride)

PRODUCT DESCRIPTION

Yellow colour powder, which contains 50% of Chlortetracycline Hydrochloride (equivalent to 46.46% of Chlortetracycline), i.e. 500mg of Chlortetracycline Hydrochloride (equivalent to 464.6mg of Chlortetracycline) in 1 g.

PHARMACODYNAMICS & PHARMACOKINETICS

Chlortetracycline hydrochloride is a bacteriostatic antibiotic belonging to the tetracycline group, which acts by inhibition of protein synthesis.

Chlortetracycline hydrochloride is readily absorbed and distributed in the tissues in chickens and pigs. Blood concentrations may be enhanced by administration with citric acid.

INDICATION

For use in the treatment of infections due to susceptible bacteria in broiler chickens and pigs.

Chickens: The product is used in the treatment and control of colibacillosis secondary to infectious bursal disease, chronic respiratory disease caused by Mycoplasma gallisepticum and Escherichia coli, and Salmonella enteritidis, Salmonella typhimurium and Pasteurella multocida infections.

Pigs: The product is used in the treatment and control of respiratory diseases associated with Mycoplasma hyopneumoniae, Streptococcus suis and toxigenic strains of Pasteurella multocida. It is also used in the treatment and control of rhinitis due to Bordetella bronchiseptica and streptococcal meningitis due to Streptococcus suis type II.

TARGETED SPECIES

Broiler chickens and pigs.

RECOMMENDED DOSAGE AND ADMINISTRATION

The product is recommended for oral administration in drinking water.

Chickens: 200g of product should be dissolved in 600L of drinking water and administered daily for up to 5 days. This will provide a daily dosage of 20 - 50mg Chlortetracycline HCl/kg depending on water consumption. Dosage may be adjusted to up to 60mg/kg depending upon the severity of infection, by careful calculation of the total bodyweight of the birds and dissolving the requisite amount of product in the quantity of water consumed within 24 hours.

Pigs: In order to achieve the recommended dosage rate of 20mg Chlortetracycline HCl/kg, 200g of product should be dissolved in 500L of water and administered daily for up to 5 days. This will provide medication for 5000kg of pigs. Allowance for wastage of medicated water and reduced water intake should be made and the necessary adjustment made to the amount of product used.

Medicated water should be prepared daily and any unused water should be discarded safely. Chickens and pigs should have access only to medicated water during treatment.

CONTRAINDICATIONS

Absorption of chlortetracycline from the alimentary tract is reduced by calcium, iron, magnesium and zinc salts.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals:

After opening the sachet of a 100g or 1000g pack, contents must be used within 24 hours. Any medicated water, which is not consumed within 24 hours, should be discarded safely.

Chickens and pigs should have access only to medicated water during treatment.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Avoid inhalation of dust by wearing a protective mask. Wear either disposable half-mask respirator or a non-disposable respirator with a filter.

In case of accidental spillage on to the skin or eyes, wash the affected area with clean running water immediately. Seek medical attention if irritation occurs/persists. Wash hands thoroughly after using this product.

INTERACTIONS WITH OTHER MEDICATIONS

Absorption of chlortetracycline from the alimentary tract is reduced by calcium, iron, magnesium and zinc salts.

PREGNANCY AND LACTATION

Do not administer TP-CTC HCl 500 mg/g Powder to pregnant sows.

ADVERSE EFFECTS

Not applicable.

OVERDOSE & TREATMENT

No specific antidote is available.

WITHDRAWAL PERIOD

Chickens: 3 days.

Pigs: 6 day.

Not for use in laying birds producing eggs for human consumption.

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Target Tissues (all food producing species)	Maximum Residue Limits (MRLs) in µg/kg
Chlortetracycline	Kidney	600
	Liver	300
	Muscle	100
	Eggs	200

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

500gm

1000gm

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised August 2019