

## **Farmcare 20% w/w Chlortetracycline Hydrochloride Water Soluble Powder**

Each kg contains Chlortetracycline HCl.....200g

### **Product Description**

Light yellow powder

### **Pharmacodynamics**

Chlortetracycline hydrochloride is a bacteriostatic antibiotic belonging to the tetracycline group, which acts by inhibition of protein synthesis.

### **Pharmacokinetics**

Chlortetracycline hydrochloride is readily absorbed and distributed in the tissues in chickens and pigs. Blood concentrations may be enhanced by administration with citric acid.

### **Indication**

For use in the treatment of infections due to susceptible bacteria in broiler chickens and pigs.

Chickens: Farmcare 20% is used in the treatment and control of colibacillosis secondary to infectious bursal disease, chronic respiratory disease caused by *Mycoplasma gallisepticum* and *Escherichia coli*, and *Salmonella enteridis*, *Salmonella typhimurium* and *Pasteurella multocida* infections.

Pigs: Farmcare 20% is used in the treatment and control of respiratory diseases associated with *Mycoplasma hyopneumoniae*, *Streptococcus suis* and toxigenic strains of *Pasteurella multocida*. It is also used in the treatment and control of rhinitis due to *Bordetella bronchiseptica* and streptococcal meningitis due to *Streptococcus suis* type II.

## **Recommended Dose**

Farmcare 20% is recommended for oral administration in drinking water.

Chickens: 500g of product should be dissolved in 600L of drinking water and administered daily for up to 5 days. This will provide a daily dosage of 20-50mg Chlortetracycline HCl/kg depending on water consumption. Dosage may be adjusted to up to 60mg/kg depending upon the severity of infection, by careful calculation of the total bodyweight of the birds and dissolving the requisite amount of Farmcare 20% in the quantity of water consumed within 9 hours.

Pigs: In order to achieve the recommended dosage rate of 20mg Chlortetracycline HCl/kg, 500g of product should be dissolved in 500L of water and administered daily for up to 5 days. This will provide medication for 5000kg of pigs. Allowance for wastage of medicated water and reduced water intake should be made and the necessary adjustment made to the amount of product used.

Medicated water should be prepared daily and any unused water should be discarded safely.

Chickens and pigs should have access only to medicated water during treatment.

## **Contraindication**

None Known

## **Warning and Precautions**

### **i. Special precautions for use in animals**

Seal after use and keep container tightly closed. Any medicated water, which is not consumed within 9 hours, should be discarded safely.

Chickens and pigs should have access only to medicated water during Treatment

### **ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Avoid inhalation of dust by wearing a protective mask. Wear either disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143.

In case of accidental spillage on to the skin or eyes, wash the affected area with clean running water immediately. Seek medical attention if irritation occurs/persists. Wash hands thoroughly after using this product.

### **Interaction of Other Medicaments**

Absorption of chlortetracycline from the alimentary tract is reduced by calcium, iron, magnesium and zinc salts.

### **Pregnancy and Lactation**

Do not administer Farmcare 20% to pregnant sows

### **Side Effects**

Not applicable

### **Symptoms and Treatment of Overdose**

No specific antidote is available.

### **Storage Condition**

Store at 30°C, store in dry place, and protect from sunlight.

Keep out of reach of children/Jauhkan dari kanak-kanak.

### **Shelf life**

Shelf life (as packaged for sale): 24 months from the date of manufactured

Shelf life (after first opening): 24 hours

Shelf life (after reconstitution or dilution): Any medicated water that is not consumed within 9 hours should be discarded safely.

### **Withdrawal Period**

Chickens: 3 days

Pigs : 6 days

Not for use in laying birds producing eggs for human consumption.

### **Packing**

Aluminium foil of 1kg

### **Product Registration Holder**

Peterlabs Sdn Bhd.

Lot 16014 (PT No. 24341),

Jalan Nilam 3, Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan, Malaysia.

### **Manufacturer**

Pucheng Chia Tai Biochemistry Co., Ltd.

No. 56 Dashixi, Putan Village, Pucheng,

Fujian 353400 P.R. China.

### **Date of revision of package insert**

25 July, 2019