

VP AMOXCILN 50% W/W WATER SOLUBLE

STRENGTH OF ACTIVE SUBSTANCE

COMPOSITION: per 1gm

Amoxicillin Trihydrate 500mg

DESCRIPTION:

Appearance: A free flowing almost white powder with typical Amoxicillin odour.

(Specification: A white to almost white free flowing powder with typical Amoxicillin odour.)

PHARMACODYNAMICS:

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against Gram positive and Gram negative bacteria. IT owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

PHARMACOKINETICS:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids.

Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals.

Biotransformation appeared a more important route of elimination in birds than in mammals.

TARGET SPECIES:

Poultry (chicken,ducks,turkeys);Swine

INDICATIONS:

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

Not effective against beta-lactamase producing organisms.

Use of the product should be based on susceptibility testing and it should take into account official and local antimicrobial policies.

Pigs: For the treatment of salmonellosis and pasteurellosis.

RECOMMENDED DOSAGE:**Chickens:**

The recommended dosage is 15 mg amoxicillin trihydrate per kg bodyweight

(equivalent to 30mg of VP Amoxciln 50% w/w water soluble powder).

The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks:

Recommended dosage is 20 mg amoxicillin trihydrate/kg bodyweight

(equivalent to 40mg of VP Amoxciln 50%w/w water soluble powder

for 3 consecutive days.

Turkeys:

Recommended dosage is 15-20 mg amoxicillin trihydrate/kg bodyweight (equivalent to 30mg -40 mg of VP Amoxciln 50% w/w water soluble powder) for 3 days or in severe cases for 5 days

Pigs:

The recommended dosage is 20 mg amoxicillin trihydrate per kg bodyweight

(equivalent to 40mg of VP Amoxciln 50%w/w water soluble powder) for 5 day via drinking water

Oral administration in drinking water

Administer in the drinking water to give 20mg/kg bodyweightdaily

(equivalent to 40mg of VP Amoxciln 50% w/w water soluble powder). The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days.

For administration via the drinking water, the solution should be prepared with fresh portable water.

Any medicated water which is not consumed within 12 hours should be discarded.

In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst on treatment.

ROUTE OF ADMINISTRATION:

Oral

CONTRA-INDICATIONS/SAFETY DATA:

VP Amoxiclin 50% W/W Water soluble powder should not be administered to rabbits, hamsters, gerbils and guinea pigs, or to birds producing eggs intended for human consumption.

WARNING AND PRECAUTIONS:

WARNING & PRECAUTION

For Veterinary Use Only"

Keep Out of the Reach of Children

Jauhi dari Kanak-Kanak

Special warnings for each target species

None.

Special precautions for use

Special precautions for use in animals

Pigs:

The uptake of medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed or water, animals should be treated parenterally. Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa.

Allergic reactions to these substances may occasionally be serious.

Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.

Handle this product with great care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non disposable respirator to European Standard EN140 with a filter to EN143

Wear gloves during preparation and administration of medicated water or liquid feed

Wash any exposed skin after handling the product or medicated water or feed

Wash hands after use

INTERACTIONS WITH OTHER MEDICAMENTS:

None known.

PREGNANCY AND LACTATION:

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin. Use only according to the benefit/risk assessment of the responsible veterinarian

SIDE EFFECTS:

None known

SYMPTOMS AND TREATMENTS OVERDOSE:

No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

WITHDRAWAL TIMES:

Chickens (meat & offal):	1 day
Ducks (meat & offal):	9 days
Turkeys (meat & offal):	5 days
Pigs (meat & offal):	2 days

Not for use in laying birds producing eggs for human consumption

STORAGE CONDITION:

Store Below 30 °C

Protect from light and moisture

DISPOSABLE :

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements

Shelf Life (as packaged for sale)

2 years as packaged for sale

Shelf Life (after first opening of container)

This product should be discarded after 12 hours of opening of the container

Shelf life (after reconstitution or dilution)

This solution must be discarded after 12 hours after reconstitution or dilution

PACKING:

1 kg powder packet in aluminium sachet

MANUFACTURED BY:

Hangzhou An-I-Mate Animal Pharmaceutical Co.,Ltd

No.2 ,Fengdu Road,
Fengdu Industrial Garden
Pingyao, Yuhang, Hangzhou,
China

DISTRIBUTOR:

Yeam Heng Trading Co.
1770 Kulim Road 14000 BUKIT MERTAJAM
PULAU PINANG MALAYSIA

DATE OF REVISION OF PACKAGE INSERT:

SEPTEMBER 2019