

K-WORM CHOC CHEWTAB**Active Ingredient:**

Each tablet contains -
Albendazole 200mg

Presentation:

Box of 2's

Product Descriptions:

Brown coloured, chocolate flavoured, round and flat uncoated tablet.

Pharmacology:

Albendazole exhibits vermifugal, ovicidal and larvacidal activities. It inhibits tubulin polymerization in anthelmintics and thus depletes their energy level until they become inadequate for survival.

Albendazole is poorly absorbed from the gastrointestinal tract, but rapidly undergoes extensive first - pass metabolism. The main metabolite, albendazole sulfoxide, which is widely distributed in the body especially in the bile and the CSF, has anthelmintic activity and have a plasma half-life of 8.5 hours. It is approximately 70% protein-bound to the plasma protein. It is then mainly eliminated in the bile while a small amount is excreted in the urine.

Indications:

For the treatment of single or mixed infestations of intestinal parasites.

Dosage and Administration:

For pinworm, whipworm, roundworm and hookworm infestation:

Adults & children above 2 years:

Chew 2 tablets as a single dose.

Children 1 to 2 years:

Chew 1 tablet as a single dose.

For tapeworm and threadworm infestation:

Adults & children above 2 years:

Chew 2 tablets once daily for 3 consecutive days.

Symptoms and Treatment of Overdose:

Symptoms: Gastrointestinal symptoms may occur in Albendazole overdose such as nausea, vomiting, diarrhoea and abdominal pain.

Treatment: Gastric lavage may be performed in first 2 to 3 hours after ingestion and other supportive and symptomatic treatment may be carried out if necessary.

Contraindications:

Person who exhibits hypersensitivity to the drug or other benzimidazole derivatives. Should not be administered during confirmed or suspected pregnancy.

Precautions:

All family members and others in close contact should be treated at the same time to prevent reinfestations.

If infestations persist or reoccur, please consult a doctor.

Not recommended for children under 1 year unless advised by a doctor.

Unsuitable for phenylketonurics.

Patients treated for neurocysticercosis should receive appropriate steroid and anticonvulsant therapy to minimize cerebral hypertensive episodes.

Use in Pregnancy and Lactation:

Should not be administered during confirmed or suspected pregnancy.

Use with caution during lactation as extent of excretion in breast milk is not known.

Undesirable Effects:

Albendazole is generally well tolerated. Symptoms reported include gastrointestinal symptoms and headache.

Interactions with Other Medicament:

Caution should be taken when Albendazole is administered concurrently with these medications.

Coadministration of Carbamazepine may decrease the efficacy of Albendazole.

Concomitant use with Cimetidine, Dexamethasone and Praziquantel, the plasma levels of the active metabolite of Albendazole may increase.

Storage:

Store below 30°C. Protect from light and moisture.

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak.

Shelf Life:

3 years from the date of manufacture.

Registration No.: MAL19126029XCZ

X.TPwWC-L.01

Date of Revision: 19 January 2023

Product Registration Holder

Powerlife (M) Sdn Bhd

No.8-1, Jalan PDS 4,

PD Sunggala Gateway,

Telok Kemang,

71050 Port Dickson,

Manufacturer

WINWA MEDICAL SDN. BHD.

1952, Lorong I.K.S. Bukit Minyak 3,

Taman I.K.S. Bukit Minyak,

14000 Bukit Mertajam,

Pulau Pinang, Malaysia.

Size: 75mm x 100mm

 Black